

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/



Dr. Laurence Welti Gift

alkaline Metals Sodium ammonim Lithium Potassium esto are more racity absorbed better divisties there sodium call Sodium salts are > deforteent a Letter purgatives. announce is a general atimular magnesia + mages recemble Kalla es Ulum a ca salto are mild act Ulum is an emptie hime water is autematic. Ba al mould pharmacologically all the Pt altrare local redatives ag. more powerfully cousin the For Cle is a powerful cometie.



Potassium eath are more rasily absorbed better directies than sodium salt Sordinum entre are > deposeent a better purgatives, annument is a general atimular magnesia + mages recemble Kalle e alum a ca calto are mild agt Ulum is an emetic hims water is autematic. BA al would pharma cologically of the Pr alts are local sedatives ag. more powerfully sanche the For Cle is a powerful constre. C-50 4

exter analdehyde .- not depressant sulphonal - assont depress heart. lamplene hydrate - not depressant. Joinel antipyretic + analges autamilia Pheracitin - Slightly Exalgin Phenosoll hydrochloude hallin sulphate -

antisepton

	1
Carbolia and = Phunol	4-19.
cureti	2-2m
quained 60-90% errorate	2-10 -
enerate carbonale 92 % "	15-60
quained Burgoute	2 - 10 "
carbonate	4" - 801-
" sainfate	4 30
creclin	1-5-
fugast "	
Jodeform - alone 91% I	1-392.
anistre 45 of I	
3-de 88 % I	- 3 pc
Europhen 27 % I	
Naphtalin Jantes autisipais asspert	
naplited Interactionperio	1-200
assper	1 - 20 4
Reconcin	75-60.
	2 - 5°,
Formaldehyde	
Sacham - not used as anti	
mithydene blue	• -

Drugo acting on Resp. Organo Squiel sinega Sauguinaria preac Sobelia Vigitalia group. acouste group Digitalia assiste Veratrum 5quill 3 thophanthus veratring Convallaria Seoparius cactus grandiflorus adomidin aposymum atropine group Salicylia growt Salisin Balicylie te. 6.60 Agosinamore na Salicylete 5- 60 Hyoseine Haturine Sithinm Salington 1-15 Scopolamine Balol (Phenyl ..) 5-30 Duboisine methyl Salicylate 1-5 m Oleman gunttherine >5 .. Betulae 1-5 Balophen 5-15 hay sal. 4gr

A SYLLABUS

OF

MATERIA MEDICA

COMPILED BY

WARREN COLEMAN, M.D.

Instructor in Clinical Medicine and Materia Medica in Cornell University, Medical Department; Visiting Physician to City (Charity) Hospital; Assistant Visiting Physician to Bellevue Hospital



NEW YORK
WILLIAM WOOD & COMPANY
1899



and Purdiana doplyllum

COPYRIGHT WILLIAM WOOD & COMPANY 1898



PRESS OF STETTINER BROS., 22, 24 & 26 READE STREE NEW YORK. 1899

PREFACE.

An experience of several years in actual class-room work has convinced the author that Materia Medica is one of, if not quite, the most difficult subject to acquire in the field of medicine. Its study from the medical standpoint cannot rest upon a rational basis until the student is brought face to face with the patient at the bedside. Except in its application, Materia Medica is solely a matter of memory and must be learned and relearned many times Accordingly the attempt is here made to assist the memory as much as possible by condensing the facts, repeating the doses, and by grouping the drugs in various ways.

Some of the more unimportant drugs have been omitted. It is to be regretted that the pruning process could not have been more radical.

It is specially intended that the present volume shall not take the place of, but be supplementary to, text books.

The doses adopted have been taken from White's "Materia Medica and Therapeutics." The author wishes to acknowledge his indebtedness to the same volume for the majority of facts here incorporated, and to Messrs. William Wood & Co. for their courtesy and liberality in resetting much of the matter in order to determine upon the form best suited to the text.


5 WEST THIRTIETH STREET. November 1st, 1898.

TABLE OF CONTENTS.

PAGES	
	I.—SCHEMA FOR STUDYING THE ACTIONS
1- 3	AND USES OF DRUGS
	II.—SYLLABUS OF THE ACTIONS AND USES
4- 93	OF DRUGS
94-111	II.—The more important preparations,
	IVCLASSIFICATIONS OF DRUGS, BASED
	CHIEFLY ON PHYSIOLOGICAL AC-
112-140	TIONS
•	V.—VARIOUS GROUPINGS
141	1Drugs which may produce eruptions
141	2.—Drugs excreted in the milk
•	3.—Drugs which affect the color of the
142	feces
•	4Drugs which affect the color of the
143	urine
-13	5.—Drugs which may cause the urine
143	to reduce Fehling's solution
	6.—Gastro-intestinal irritants
146-151	7.—Alkaloids: their sources and salts

TABLE OF CONTENTS,

	PAGES
8.—Active principles of drugs other than	
alkaloids	152-156
9.—Drugs with a maximum dose of $\frac{1}{10}$	
grain or less	157-158
10.—Drugs known by common and other	
names	159-166
11.—Dosage in children	167
12.—Rules for incompatibility	167
13.—An easy method of writing prescrip-	
tions	168-169



SCHEMA FOR STUDYING THE ACTIONS AND USES OF DRUGS.

 External. (Chiefly local or direct.) Including nose, mouth, pharynx, bronchial tubes (inhalation), urethra, vagina, rectum.

II. Internal. -

A.—Alimentary tract (action local). Mouth, stomach, intestines.

B.—Circulation.

- Blood (entrance into; changes effected in; indirect effect on metabolism, indicated by urea, carbon dioxide, and temperature).
- (2) Heart (force, frequency, regularity; action direct or through nerves; blood-pressure).
- (3) Blood-vessels (effect on calibre; action direct or indirect; blood-pressure).

C.—Nervous system.

- (1) Brain (vascular changes in; action on motor or sensory areas; on special centres, cardiac, respiratory, vaso-motor, vomiting, heat).
- (2) Cord (reflex activity, anterior cornua, afferent and efferent tracts).
- (3) Peripheral nerves (afferent and efferent).

D.—Secretions.

- Saliva, gastric, pancreatic, and intestinal juices. Also physical and chemical changes in feces.
- (2) Bile (direct or indirect action).
- (3) Urine (quantity, chemical and physical changes, including amount of urea and odor).
- (4) Sweat (quantity, effect on temperature).
- (5) Bronchial (increased or diminished; if disinfected).
- (6) Milk (increased or diminished; if excreted in).

E.—Special actions.

- (I) Eye (pupil, accommodation, tension).
- (2) Genito-urinary tract.
- (3) Muscles (voluntary, involuntary).
- (a) Does it easily produce symptoms of poisoning?1. Children. 2. Adults. 3. Tolerance.

(b) What symptoms?

Anorexia, nausea, vomiting, purging; cardiac; respiratory; nervous; tinnitus aurium; renal congestion; eruptions.

(c) How administered?

- 1. With reference to meals.
- 2. Night or morning.
- 3. Diluted or undiluted.
- 4. Flavored.
- 5. In capsules or wafers.

n. acetanilid pour der (acetanilid) 2002 /2 31

SYLLABUS OF MATERIA MEDICA.

ABSINTHUM (wormwood)	8–40 gr.
Use. Stomachic.	
ACACIA (gum Arabic): Mucilago a	-
Uses. Demulcent, vehicle.	
Actions. Cardiac depressant. V. Antipyretic. Analgesic. Uses. Chiefly as antipyretic and analg ACIDUM ACETICUM: Ac. aceticum dil., 6%	so-constrictor. Muralgae esic. Dispunioration Scratica Transcription Market Constraints
ACIDUM ACETICUM GLACIALE	_
antipying Salipying Sodopying Phonosition Exalging (southap) acutamilia (so to phonosity)	0-20 gr. 6-30. 2-16. 5-10. 4-3 10-16-

THE ACTIONS AND USES OF DRUGS. 5
Syr. acidi citrici
ACIDUM HYDROCHLORICUM:
Ac. hydrochloricum dil., 31% 5-20 ¶.
ACIDUM LACTICUM.
ACIDUM NITRICUM:
Ac. nitricum dil., 10% 5-30 M. " nitrohydrochloricum.
и dil 5-30 щ,
ACIDUM PHOSPHORICUM:
Ac. phosphoricum dil., 10% 5-30 M.
ACIDUM SULPHURICUM:
Ac. sulphuricum dil., 10%10-30 M.
" ceini of within 5-15 M.
ACIDUM TARTARICUM 8-30 gr.
Actions. Ext. Irritants, except citric. Caustics, Astringents. Hemostatics. Refrigerants when dilute. Int. Sialagogues. Before meals check gastric juice. Sulphuric astringent in intestine. Cholagogues, especially nitric and nitrohydrochlo- ric, Acetic, citric, and tartaric increase alkalinity of blood, others diminish it.
, · · · · · · · · · · · · · · · · · · ·

sile duct.

Uses. Ext. To destroy warts, condylomata, etc. Hemostatics. Refrigerant in fevers. Int. To allay thirst. Dyspepsia. Sulphuric as hemostatic in stomach and intestines, and in diarrhoea. Cholagogue. Vegetable acids for scurvy, and in combination as diuretics and in gout. Sulphuric as anhidrotic in tuberculosis.

ACIDUM BORICUM..... 5-15 gr. Glyceritum boroglycerini. Sodii boras (borax) Actions. Local disinfectant and antiseptic. Uses. Wounds, ulcers, etc. Vaginitis, urethritis, etc. Aphthous stomatitis. Aciduri CARBOLICUM (phenol) 15-2011 - 1 gr. Ung. acidi carbolici, 5%, Glyceritum acidi carbolici, 20%.......... 2-5 III. Actions. Ext. Antiseptic, disinfectant, deodorant, parasiticide. Irritant, caustic. Local anæsthetic. Int. Small doses, antiseptic. Large doses, general depressant. Urine dark from pyrocatechin and hydrochinon. Actions. Said to diminish secretions of mucous surfaces. Uses. Chronic bronchitis. Pyelitis, etc. Liquor antiseptions = (Borie acid 27), Bangoie : Thymole alcohol is the antidote for corbs

	THE ACTIONS AND USES OF DRUGS. 7
1	ACIDUM HYDROBROMICUM DILUTUM, 1-1 dr.
	The state of the s
	Actions. Int. Like the bromides. Uses. Int. To relieve noises in the ears. To prevent cinchonism. To dissolve quinine.
	ACIDUM HYDROCYANICUM DILUTUM (prussic acid)
	Actions. Ext. Sedative, local anæsthetic. Int. Sedative. Cardiac, vaso-motor, respiratory depressant. Uses. Itching. Gastric pain, vomiting. For coughs.
	ACIDUM*OLEICUM.
	Use. Solvent of drugs for inunction.
	ACIDUM STEARICUM.
	Uses. Making glycerine suppositories. In combina- tion with zinc and copper for skin diseases.
	ACIDUM SULPHUROSUM 1-2 dr.
	Actions. Ext. Disinfectant, deodorant, parasiticide. Int. Disinfectant in stomach and intestine (?). Uses. In form of gas from burned sulphur to disinfect rooms, etc. Ringworm and foul sores,
	ACIDUM TANNICUM 1-20 gr.
	Collodium stypticum 20%
	Troch, ac. tan. (1 gr.)
	Ung. ac. tan 20%
	Glyceritum ac. tan 20%

Actions. Ext. Astringent. Hemostatic. Int. Astringent and hemostatic in stomach. Changed to gallic acid in intestine. Uses. Ext. As general astringent and hemostatic. Int. In other drugs, as catechu, for diarrhoea.
ACONITUM (monkshood). \(\) \(
Part - W.W. W. M. T. 2 - 81.
Est a
Ext. a. fl ½-2 m.
Tr. a 3. 5 1-5 m.
(Fleming's tr. twice as strong.) 12-2m.
Aconitina. 2670.#.20.) 100 gr. with caution.
Napellina ⅓ gr
Actions. Ext. Tingling, numbness, local anæsthesia. Int. Heart slowed, force lessened. Blood-pressure falls. Vessels dilate (?). Depresses nerves of sen- sation, temperature, and later motion. Pain re- lieved. Antipyretic. Diaphoretic. Diuretic (?).
Uses. Ext. Neuralgia, etc. Int. Neuralgia. Febrile
Jungiseases. Tonsillitis, laryngitis, etc.
At. for 50 lb. child. 3 doses enough
ADEPS (lard):
Adeps benzoinatus.
Ceratum.
" resinæ.
Unguentum,
Oleum adipis.
lices Emollient Making ointments

Uses. Emollient. Making ointments.

THE ACTIONS AND USES OF DRUGS. deps Lanae ADEPS LANÆ HYDROSUS (lanolin): Uses. Basis for inunctions. Europe + aria ADONIDIN (plucoside fr. adonis vernalis) ... 1-1 gr. Actions. Cardiac stimulant. Slows heart. Raises blood-pressure, which subsequently falls from vasomotor paralysis. Action prompt. Wat comme Uses. Like digitalis. ÆTHER 5-60 M. Spr. ætheris..... 1-1 dr. Oleum sethereum. (therese wil) Spr. ætheris co. (Hoffman's anodyne) ... 5-60 m. Actions. Ext. Evaporating, produces cold and local anæsthesia. Confined, irritant. Int. Like alcohol and chloroform. Stomachie, carminative. Cardiac stimulant. General anæsthetic Respiratory stimulant. Outies andie Uses. Ext. Small operations. Int. As stomachic. Flatulent dyspensia. Fainting, heart failure, pulpitation, Antispasmodic, asthma, General anasthetic. Cautions. Do not use in: 1. Bronchitis or pulmonary disease. 2. Renal disease. 3. Heart disease. 4. Aneurism. 5. Arterial sclerosis. 6. In hot climates. 7. About artificial light. ÆTHER ACETICUS.....20-60 M. Actions and Uses. Stimulant, antispasmodic, carminative. an alle laidal stone + 1 compai, asthme

ÆTHYLENE BICHLORIDUM (Dutch liquid):

Action and Use. General anæsthetic.

ÆTHYL BROMIDUM (hydrobromic ether):

Actions. Rapid, transient anæsthetic, without total loss of consciousness.

Uses. Minor surgery.

Use. Anhidrotic in night sweats.

*ALCOHOL

Alcohol absolutum	99%
deodoratum	92.5%
**	91%
Spr. frumenti (whiskey)	44-50%
" vini gallici (brandy)	39-47%
Gin, rum, liqueurs	
Alcohol dilutum	41%
Vinum album fortius	20-25%
" portense (port)	15-25%
" xericum (sherry)	15-20%
Madeira	15-20%
Champagne	10-13%
Burgundy	10-13%
in about	

traves tody unchanged invitation

give teretures in nephritis.

Vinum album
" rubrum 10-14%
Claret 8-12%
Cider
Ale or stout 5- 9%
Beer or porter 2- 5%
Kumyss
Actions. Ext. Disinfectant. Refrigerant. Astringent. Anhidrotic. Hardens skin. When confined, rubefacient. Int. Stomachic. Local anaesthetic. Diminishes oxidation. Antipyretic
Cardiac stimulant; after-effect, exhaustion. Vaso-dilator. Raises blood-pressure. Diaphoret- ic. Diuretic. Small doses, nervous stimulants;
large, narcotic. Uses. Ext. Refrigerant, anhidrotic. Headache.
Prevent bedsores Int Aid direction Vomit-
ing. Diarrhœa, Cardiac stimulant, Insomnia.
Diuretic. Carbolic 7 occomen -
ALLIUM (partie) a sute heart failure in Pries. etc.
ing. Diarrhoea. Cardiac stimulant. Insomnia. Diuretic. Cardiac stimulant. Insomnia. Diuretic. Cardiac stimulant. Insomnia.
Actions. Stomachic. Cardiac stimulant. Expecto-
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspensia.
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspensia.
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Book along to gr.
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspensia.
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Book along to gr.
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Deal along 10 gr. SOCOTRINA E. Aprica 1-10 gr. Ext. a. 1-3 gr. A. purificata. 1-10 gr.
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Book along 10 gr. 50COTRINA. C. along. 1-10 gr. Ext. a
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Barbally—10 gr. SOCOTRINA & Africa 1-10 gr. Ext. a
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Barbally—10 gr. SOCOTRINA & Africa 1-10 gr. Ext. a
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Barbales 10 gr. SOCOTRINA & Africa 1-10 gr. Ext. a. 1-3 gr. A. purificata. 1-10 gr. Reschol in along dozes of peptitons & popular to the period of
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Barbally 10 gr. "SOCOTRINA & Africa 1-10 gr. Ext. a. 1-3 gr. A. purificata. 1-10 gr. Parally done of peptloms & pept
Actions. Stomachic. Cardiac stimulant. Expectorant. Uses. Dyspepsia. ALOE BARBADENSIS. Barbales 10 gr. SOCOTRINA & Africa 1-10 gr. Ext. a. 1-3 gr. A. purificata. 1-10 gr. Reschol in along dozes of peptitons & popular to the period of

12 SYLLABUS OF MATERIA MEDICA.

	Pil. a., gr. 2 1-4 pills.
n.alm-	Pil. a, et asafœtidæ, gr. 1] 1-5"
S04 00	\ ←Pil. a. et ferri, gr. 1
	Pil, a. et mastiches (" Lady Webster "),
	gr. 2 I-3 "
	Pil a. et myrrhæ, gr. 2 2-10 "
nalow 10	Tr a
iqueria "	Tr. a. et myrrhæ lligin Proprietatie 1-24 dr.
iquoria ros	Aloinum
	Aloinum. 1-24 dr. Remarket dr. 1-24 dr. Aloinum. 1-2 gr. 2
	Actions. Stomachic. Purgative-15-20 hrs. Acts
	chiefly on colon. Gripes. Cholagogue. Em-
	menagogue. Wented by wilk
	Uses. Stomachic, Habitual constipation. Chlorosis and anæmia with iron. Amenorrhea.
	old and anicollies with months a simple of the simple of t
	ALTHÆA (marshmallow):
	XXX
	My man
	Uses. Demulcent Flavoring agent.
	t vianaulani
	ALUMINUM:
	Alumen 5–30 gr.
	Alumen exsiccatum.
	A. hydras 1-10 gr.
	A. sulphas.
	Actions. Ext. Coagulates albumin of sores and ul-
	cers, Astringent, Hemostatic Int. Astringent,
	mouth, stomach, intestines. Direct emetic.

Uses. Ext. Destroys unhealthy granulations. Eczema, leucorrhœa, gleet, conjunctivitis, etc. Piles, cuts, etc. Int. Tonsillitis, etc. Emetic. Gastric hemorrhage Diarrhœa.

AMMONIUM

Aqua amme	oniæ fortior Well diluted.	3-6	η.
Spiritus am	moniæ	10-60	η.
Aqua amm	oniæ		η.
	n ammoniæ.	4 9	n
	Rubefacient, vesi erally in stomach.		
ratory stimi	ulant. Raises bloo	od-pressure.	

Uses. Ext. Counter-irritant. Insect bites. Syncope (inhalation). Int. Dyspepsia. Cardiac and respiratory stimulant in collapse. Snake-bites (hypodermically).

A. carbonas ... 2-15 gr.

Spiritus ammoniæ aromaticus (sal volatile). 4-2 dr.

Actions and uses. Gastric stimulant, Carminative.
Cardiac and general stimulant. Expectorant.
Emetic.

Uses. Ext. Chr. pharyngitis, otitis media, laryngitis, bronchitis (by inhalation). Int. Expectorant.

MH 1) CO3 3.4 90

Liquor a.	acetatis	(spirit	of Mindererus)	2-8 dr.
-----------	----------	---------	----------------	---------

Actions	and	uses.	Diaphoretic	and	diuretic	in
febril	le con	ditions	and nephritis.			

	febrile conditions and nephritis.
	AMYGDALA (almond):
	A, amara (bitter almond):
	Oleum a. a 1-1 M.
	Aqua a, a 2-4 dr.
	Spr. a, a15-45 M.
	A. dulcis (sweet almond):
	Emul. amygdalæindeterminate.
	Syr. " "
	Ol. amygdalæ expressum (fr. bitter or sweet)
	Actions and uses. Bitter almond for vehicle and flavoring. Sweet almond, demulcent, vehicle, and to replace flour in diabetes.
	AMYLENI HYDRAS 1-1 dr.
	Action and use. Hypnotic. Safe.
*	AMYLINITRISdose. { inhaled 1-3 M. internally \frac{1}{2}-1 M.
100	Actions. Int. Increases cardiac frequency—vagus inhibition. Vaso-dilator. Symptoms of cerebral
hitrita	Struck withite (Sweet Sp. Mitter)
tratio	general trinituate (Vitta gly)

THE ACTIONS AND USES OF DRUGS.

congestion. Respiratory stimulant. Antipyretic, Depresses reflexes.

Uses. Cardiac and aneurismal pain. Angina pectoris. Epilepsy. Migraine. Tetanus, strychnia poisoning. Asthma. Whooping-cough.

Glyceritum amyli.

Uses. Chiefly for mechanical properties.

ANISUM (anise)	10-30 gr.
Oleum a	1-5 TL.
Aqua a	1-I oz.
Spr. a	1-2 dr.

Actions and uses. Of volatile oils.

ANTHEMIS (chamomile)..... ½-2 dr.

Action and use. Stomachic bitter.

OLEUM ANTHEMIDIS..... 1-4 M.

Actions and uses. Like other volatile oils.

ANTIKAMNIA (proprietary):

Said to contain acetanilid, sodium bicarbonate, caffeine, and tartaric acid.

ANTIMONIUM:
A. sulphidum.
A. sulphidum. " purificatum. A. sulphuratum. A. sulphuratum. 1-1 g A. oxidum. 1-4 g Pulvis antimonialis (James powder), 3-15 g
A. sulphuratum Revery mess 1-1 g
A. oxidum
Pulvis antimonialis (James' powder), 3-15 gr
A. et potassii tartras (tartar emetic):
(diaphoretic 10 1 gr
diaphoretic $\frac{1}{10}$ gradiac depressant $\frac{1}{2}$ -1 gr
emetic I-2 co
Syr. scillæ co. (hive syrup). 2.76. 5-60 III Caphos Caphos Caphos Actions. Ext. Irritant. Tartar emetic produce
Syr. scillæ co. (hive syrup) 2.% 5-30 11
~ Calebra
local pustular eruption. Int. Irritant Direct and indirect emetic. Cardiac depressant. Powerful nerve depressant. Large doses antipyretic Expectorant. Diaphoretic. Cholagogue. Uses. Ext. Rarely as caustic. Int. Emetic. Cardiac depressant in febrile conditions. Bronchitis Not used as much as formerly.
OfficialANTIPYRIN 3-20 gr
Actions. Analgesic, antipyretic. Cardiac depres sant. Vaso-constrictor. Uses. Neuralgia, etc. Hyperpyrexia. Hemostati in epistaxis and hemorrhoids.
APIOL (a camphor from parsley) 5-15 gr

Uses. Antiperiodic. Dysmenorrhæa. Amenorrhæa.

Liquid apiol is an alcoholic extract of parsley seeds. Dose, 2-8 m.

Use. Diuretic.

APOMORPHINÆ HYDROCHLORAS: (45 H20)

doses subcut. 15 gr. by mouth 10 10 gr.

Actions. Indirect emetic. Stimulates bronchial secretion. Other actions which are not toxic are due to the vomiting.

Uses. Poisoning, etc. Expectorant.

1 If It is emerald grien it is no good)

AQUA HYDROGENII DIOXIDI (hydrogen peroxide) 1-3 dr.

Actions. Ext. Gives up oxygen readily and is antiseptic and disinfectant. Decomposes pus. Int. Gives oxygen to blood. Probably stimulates nervous system.

Uses. Ext. Washing wounds, fistulæ, etc. Mouth wash in diphtheria. Int. Dyspepsia.

apomorphime Helis the hydrochlar of an artificial alkaloid abtained to heating morphime in accord to with an excess of Hel. New tosses one molecule of the

ARGENTUM (silver):	
A. nitras (lunar caust	ic) +1 gr.
Nitras dilutus.	The state of the same
" fusus.	
A. oxidum	1-2 gr.
A. cyanidum.	
A iodidum	1 -1 gr.
4	
Said to have little action	ingent. Int. Nervous dis-
ARISTOLUM. = 346	guel iodidum
Action and use. Like i	
ARNICA:	
A. flores	10-20 gr.
A-radis-	5-20 gr.
The state of the s	5-10 gr.
	5-20 TL.
	5-10 M.
Empl. a.	******
Actions and uses. Lil	e volatile oils.
ouin 1.50% solin	
- 100 doctate 3	Ta
E (-ag citrate)	4000 170
got 80% a.	(soe H20)
tamme 107	To arlie.
anne 10 7	


THE ACTIONS AND USES OF DRUGS, 19
ARSENUM: arsenitriosadum
Acidum-arsenosum 1 gr.
Liquor potassii arsenitis (Fowler's
solution), 1% * H Cos 2. % 2-10 m.
Liquor aridi arsenosi, 18 (ALLS 5-10 M.
Sodii arsenas 10 gr.
Liquor sodii arsenatis (Pearson's
solution), 1% 1-10 M.
Arseni iodidum 10 gr.
Liquor arseni et hydrargyri iodidi
(Donovan's solution), āā 1% 1-10 M.
Actions. Ext. Caustic on raw surfaces. Int. Gas-
tric irritant in large doses. In small doses gastric
and duodenal stimulant. Hematinic. Alterative.
Uses. Ext. As caustic, but dangerous. Int. To de-
stroy pulps of teeth. Dyspepsia. Some cases of
womiting. *Some cases of diarrheea. Chronic
superficial skin diseases. Primary anæmias, as pernicious anæmia, leukæmia, Hodgkin's disease.
Secondary anæmias, as chlorosis, etc. Antiperiodic
and vague malarial manifestations. Rheumatoid arthritis. Chorea, Asthma variage, goods
arthritis. Chorea, Asthma. vuralque, gastrata
sphincters relax at the cometime.
ASAFŒTIDA:
Emul. a
MIXIMA 1-5 pills.
Pil. a., 3 gr
Tr. a
* love in children when the
APPENDING TO DETECT OF THE PERSON OF T

THE ACTIONS AND HERE OF DRIES

Actions.	Of volatile of	ils. Stimul	ates perist	alsis	5.
	ith purgative				
taste,	hysteria.	nalina	aren		

ASPIDIUM	(male fern, filix-m	as)	-1-1 dr.
Oleoresin	a a		1-1 dr.

Action and use. Anthelmintic for tapeworm.



Action. Increases amt. of oxygen in blood (Penzoldt).

Uses, Émphysema, chronic bronchitis, asthma.

AURANTII CORTEX (orange peel) :

A, amari cortex:	
Ext. a. a. fl	1-1 dr.
Tr. a. a	1-2 dr.
A. dulcis cortex.	
Syr. aurantii	vehicle.
Tr. " dulcis	1-2 dr.
Oleum aurantii corticis:	
Spr. aurantii	vehicle.
Spr. " co	
Elixir aromaticum	**

Actions and uses. Flavoring agent. Bitter.

AUDANTII FLORES.

AUKANIII FLUKES:
Oleum aurantii florum:
Aqua a. f. fortior indeterminate.
Syr. a. f
Actions and uses. Flavoring agent.
AURUM (gold): A. et sodii chloridum
Actions. Int. Promotes digestion. Stimulates brain. Aphrodisiac. Uses. Dyspepsia. Gastroduodenitis. Hypochondriasis. To absorb connective-tissue formations. Tertiary syphilis.
BALSAMUM PERUVIANUM 10-30 m.
Actions. Ext. Stimulant. Disinfectant. Int. Carminative. Expectorant. Uses. Ext. Wounds, sores, and chronic eczema. Pediculi, scabies, ringworm. Int. Chronic bronchitis.
BALSAMUM TOLUTANUM
Action. Like bals, of Peru. Use. Expectorant.

1	BARIUM:
[العديد	B. dioxidum.
at .	and the same of th
	B. chloridum 10 1 gr.
	Actions. Like digitalis. Stimulates muscle of intestine and bladder.
	Uses. Mitral insufficiency. Hemorrhage. Atony of intestine or bladder.
	PELLADONNA / Walter Habitat Cario
	BELLADUNNA (deadly nightshade):
	B. folia 1-5 gr.
Rad. b	Ext. b. foliorum alo 1 gr.
T.	Tr. b. " 5-30 π.
	Ung. b. "
	Empl. b. "20%.
	B. radix:
	Ext. b. radicis fl
	Lin. b.
	Atropine (alkaloid). (130 +0). 130 do gr. A. sulphas. (10.4 m). 120 do gr.
	Actions. Ext. Anodyne, local anæsthetic. Int. Vagus inhibited. First contracts, then dilates vessels. Blood-pressure rises, then falls. Deliriant. Stimulates, then depresses medullary centres. Antipyretic. Depresses all secretory nerves. Mydriatic. Paralyzes accommodation. Intra-ocular tension increased. Relaxes involuntary muscle.
	Uses. Ext. Pain. Int. Antisialagogue. With purgatives. Certain cardiac diseases. Night-sweats. Whooping-cough. Asthma. Nocturnal incontinence. Vesical or urethral spasm. With opium.
ine a	trapine before administring a
····	I amostluter to abolish raque control.
	sweating
phit	halmology
-	the mixtures

BENZOINUM: Adeps benzoinatus. Tr. b. 1-1 dr. Tr. b. co. (friar's balsam) .. Jan b-1 dr Acidum benzoicum ... 5-15 gr. Ammonii benzoas..... 5-20 gr. Actions. Ext. Antiseptic. Int. Causes appearance of hippuric acid in urine. Makes alkaline urine "acid. Disinfecting expectorant. Antipyretic. Uses. Ext. Stimulating antiseptic. Int. Bronchitis, pulmonary tuberculosis. Pyelitis and cystitis with alkaline urine. BERBERINÆ SULPHAS..... 1-10 gr. Actions and uses. Bitter stomachic. Slightly antipyretic and antiperiodic. Antiseptic. BISMUTHUM: B. subcarbonas..... 5-60 gr. B. subnitras..... 5-60 gr. B. citras..... 1-3 gr. Bismuthi et ammonii citras..... 2-5 gr. B. subgallas (dermatol) 5-10 gr. B. salicylas..... 5-20 gr. Actions. Ext. and int. Astringent. Gastric seda-Uses, Ext. As protective and astringent. Int. Nau-

sea, vomiting, gastric pain. Diarrhœa,

BORAX. See Acidum boricum.

BROMOFORM	IUM.		I-5 M.
Action and u. With care.	ses.	Anæsthetic.	Whooping-cough,
BROMUM:	set	aimed from	maca water)
		ım	
Sodii	11		
Ammonii	-66		5-30 gr.
Lithii	+6	*********	The second second
Calcii	**		15–30 gr.
Zinci	-62		1-5 gr.
Strontii			5-30 gr.
			motorial
salt. Resp minished. Uses. Int. Ep ic. Migrai stridulus. nia poisoni	Anapoilepsyine. Tetaning.	y depressants shrodisiacs. K y, especially gr Delirium trem us. Hysteria. Spermatorrhoe	except ammonium Metabolism di Britand mall Hypnot- tens. Laryngismus Neuralgial Strych- a, Nymphomania, 10-60 gr. 1-5 dr.
Action and us			thartic.
tor Jacksonia	m!	Epileps	*
ha : fa	100	n:	i.i.d.

cof	fine is aline in 1 h. execut offeet	
100	THE ACTIONS AND USES OF DRUGS. 25	. 0
100	THE STATE OF THE S	E
В	UCHU (bucco)	_
	ctions. Diuretic. Disinfectant to gu. tract. Also of volatile oils. ses. As diuretic. Pyelitis, cystitis, gonorrhæa.	
В	UTYL-CHLORAL HYDRAS (croton chloral), 5-20 gr.	0 9
A	to heart. The track of the Africa AFFEINA (theine, guaranine). 20#1.9 1-5 gr.	6 0
* C	AFFEINA (theine, sugranine) 30#30 1-5 gr.	
	C. citrata 2-8 gr.	0
	C. " effervescens 1-4 dr.	
	ctions. Cardiac schudent, increasing force and lengthening systole. Blood-pressure raised. Cer-	U
y.	ses. Aortic and mitral obstruction. Cardiac dropsy, ascites. Pleural effusion. Nephritis. Combined with cardiac depressants.	
10	acute infections, cardiac fail	in
C	ALAMUS (sweet-flag): carbial attimuted	C.
	Ext. c. fl 15-60 M.	
A	ction and use. Stomachic bitter. Carminative.	
C.	ALCIUM:	
	Creta præparata 10-60 gr.	
300	Pulvis cretæ compositus, 5-60 gr.	
- A	in apt to get tonic signif, when in infections handle with	-
me	are 2 gr. of coffine in and	4

Mistura cretæ	2-4 dr,
Hydrargyrum cum creta (gray	
powder, 38% Hg.)	1-10 gr.
TxxxXxxXxxXxxxxxxxxxxxxxxxxxxxxxxxxxxx	DE DIE
Calcii carbonas præcipitatus	5-60 gr.

Actions. Ext. Astringent. Int. Antacid, astringent.
Uses. Eczema, diarrhoca.

Maxin Case (Vienna paste).

Actions. Ext. Lime-water, astringent. Slacked lime, caustic. Int. Lime, antacid, antemetic; antidote for mineral acids, oxalic acid, zinc chloride; intestinal astringent.

Uses. Ext. Lime-water for weeping eczema. Vienna paste as caustic. Linimentum calcis for burns.

Int. Lime-water mixed with milk. Vomiting.

Diarrhea.

Calcii phosphas præcipitatus...... 5-30 gr. Syrupus calcii lactophosphatis...... 1-2 dr.

Actions and uses. Increases salts of bone. Rickets. Anæmia.

ea c	de or calactate ma he made
befo	ne operations to 4 cong of blood
1	THE ACTIONS AND USES OF DRUGS. 27
1-	Calx sulphurata 1 gr.
1	Used in suppuration.
1	Calcii ehloridum 5-20 gr.
	Used for enlarged lymph glands.
	Calcii sulphas exsiccatus (Plaster of Paris).
	Used for plaster casts.
CAI	LUMBA
	ions. Stomachic bitter. Carminative. Anthel- mintic to threadworm—injection. s. Dyspepsia. Threadworm.
CAI	MBOGIA (gamboge)
	ion. Drastic, hydragogue purgative. Gripes. In combination.
CAI	MPHORA 3-20 gr.
	Aqua c 1-2 oz.
	Lin. c. (camphorated oil). 20% Spr. c
	Ceratum c.
Act	ions. Ext. Warmth. Anæsthesia. Int. Stom-
-	phon is absorbed from munifu
~	are of symmote put now
-	the mouth.

					stimulant,	
					Cerebral	
citant	or	sedative.	Ex	pectorant.	Diaphore	etic.
Aphro	disia	C.			-	

Uses. Ext. Counter-irritant and anodyne. Int. Carminative. Coryza. Cough mixtures. Hysteria.

CAMPHORA MONOBROMATA..... 2-10 gr.

Actions and uses. Like the bromides.

CANNABIS INDICA (Indian hemp, haschisch),

	2-5	gr.
Ext. c. i	1-4	gr.
Ext. c. i. fl	2-5	Π.
Tr. c. i	5-30	TIL.

Actions. Peculiar intoxication. Deliriant. Narcotic.

Uses. Migraine, neuralgia.

CANTHARIS (Spanish fly):

Ceratum c.

Collodium cantharidatum.

Tr. c. 1-15 m.

Actions. Ext. Irritant, counter-irritant, vesicant.

Int. Gastro-intestinal irritant. Renal irritant—
albumin, blood.

Uses. Ext. Counter-irritant. Vesicant-about 8 hrs.

CARYOPI	HYLLUS (cloves)	5-30 gr.
Oleum	C	1-5 M.
	Ext. Irritant, counter-irritant, rul	
	anæsthetic. Int. Stomachic, car	
	cardiac stimulation. Stimulating secretions. Antispasmodic.	disinfect-

Actions and uses. Of volatile oils.

Oleum c..... 1-5 III.

Uses. Ext. Neuralgia. Int. Stomachic, carmina- native, antispasmodic.	
CASCARILLA 10-30 gr.	
Action and use. Stomachic bitter.	
CASSIA FISTULA (purging cassis) 1-8 dr.	Ì
CASSIA FISTULA (purging cassis) 1-8 dr.	
Use. In confection of senna.	
CATECHU. Jambur 10-30 gr.	
Tr. 9 co ½-2 dr.	
Troch. o. 1 gr	
Action and use. Ext. and Int. Astringent. Sore	
throat, diarrhœa, etc.	
OFFINA / A	
CERA (wax):	
C, flava.	

Uses. Basis for plasters, ointments, cerates.

CEREUS GRANDIFLORUS (cactus):

C. alba.

Ext. c. g. fl...... 10-30 M.

Actions. Cardiac stimulant—augmentor and intrinsic mechanism. Shortens diastole, Raises bloodpressure, Vaso-motor action not marked.

Uses. As general cardiac stimulant. Functional cardiac diseases. Aortic regurgitation. Contra-indicated in mitral stenosis.

A.M. A. Marielle 1, 1910 1946 188

THE ACTIONS AND USES OF DRUGS. 31
CERIUM:
C. oxalas 1-8 gr.
Use. Vomiting of pregnancy.
CETACEUM (spermaceti):
Ceratum c. Ung. aquæ rosæ.
Uses. Emollient. Making ointments.
CETRARIA (Iceland moss): Decoctum c
Uses. Demulcent. Food.
Oleum c 2-10 M.
Action and use. Anthelmintic for roundworm.
* CHLORAL Hydratum 5-20 gr. 7
Actions. Ext. Antiseptic. Irritant. Int. Gastric irritant. Cardiac, vaso-motor, respiratory depressant. Depresses cerebrum and cord. Hypnotic. Uses. Hypnotic. Delirium tremens, puerperal con-
whooping-cough. Hyp val = Cheral + antipprine - 15 or Character of Ch
Action and use. Hypnotic.
Eller Stormanidum 3-20
Angline hydrote & Saraldehyde

	CHLORAL FORMAMID SUM (chloralamide),
	10–30 gr.
	Action and use. Hypnotic, without after-effects.
*	CHLOROFORMUM 2-20 III.
	Aqua chloroformi
	Linimentum " 30%
	Emulsum "
	Actions. Ext. Evaporating, produces cold and local anæsthesia. Confined, irritant and vesicant. Int.
	Like alcohol. Small doses stomachic. Cardiac
	stimulant. General anæsthetic: 1st stage, stimu-
	lation; 2d stage, depression; 3d stage, reflexes abolished. No action on peripheral nerves. May
	produce vomiting.
	Uses. Ext. Chronic rheumatism and inflammations. Myalgia, Int. Toothache. Dyspepsia, Cardiac
	stimulant. General anæsthetic.
	Cautions. 1. Examine heart. 2. Remove false
has.	raised. 5. Clothing loose. 6. Stimulants at hand.
1	7. 5% chloroform, 95% air. 8. Do not begin ope-
	ration in 1st stage. 9. Watch respiration and
ع م	vagues excitability , give atropine before CHLORUM:
	CHLORUM:
	Agua chlori. Lig. chlori combos 1-4 dr. 49 Cals chlorata. Liquor soda chlorata (Labarrague's solu-
	Calx chlorata.
	Liquor sodæ chloratæ (Labarraque's solu-
Ca	ses of death
700	early operation before replices are
eiol	early operation before reflexes are
repr	casion of Heart (CHelsia a protoplamine)
La	yed poisoning dequirations
Low	ders every every to the This after CHE

Actions. Ext. Disinfectant, deodorant, irritant. Int. Rarely given. Uses. As above.
CHONDRUS (Irish moss)
CHROMIUM: Acidum chromicum.
Actions. Ext. Disinfectant, deodorant, caustic. Int. Unknown. Uses. Ext. Disinfectant for ulcers, sores, gonorrhœa, leucorrhœa, in strength of I in 40. Crystals as caustic in nares.
CHRYSAROBINUM (Goa powder, Araroba powder): Ung. c
Actions. Ext. Powerful irritant. Parasiticide. Stains yellow. Int. Gastro-intestinal irritant. Uses. Ext. Ringworm. Chronic skin diseases, e.g., psoriasis.
CIMICIFUGA (black snakeroot);
Ext, c 1-5 gr.
Ext. c. fl
Actions. Stomachic. Increases force of heart-beat. Emmenagogue. Uses. Dyspepsia. Amenorrhoza.
now very valuable.
isinfect hands:
acking powder 2 1 + 420
aff when it attings in cos + to

a a tomic in grip (4-6 gr) tostim, who has t SYLLABUS OF MATERIA MEDICA. 34 CINCHONA 10-60 gr. I-2 OZ. 5-30 gr. Ext. c. fl...... 10-60 M. Tr. c. 10 00 dr. CINCHONA RUBRA..... 10-60 gr. Tr. c. co. 20 10 1-4 dr. Quinina. (16.70 Hzb.)..... 1-20 gr. 2-10 gr. 2-10 gr. Vinum ferri amarum. 7.4.4 ettet 101-4 dr. Q. bisulphas (10. H20) 1-20 gr. Q. hydrobromadus (6: 4. 4 2.0) 1-20 gr. O. valerianas. . . . 1-30 gr. Q. sulphas.... (14.0. H.20.)..... 1-20 gr. Syr. ferri et q. et strychninæ phosphatum ... d−I dr. 1-20 gr. Q. hydrochlorashum. CB y. Mas Quinitlime sulphas...... I-20 gr. Cinchonina ... 1-30 gr. C. sulphas (6.6. H.20.)..... 1-30 gr. Cinchonidinæ sulphas. (70 H. 20.) 1-30 gr. Tr. antiperiodica (Warburg's tr.), 10 gr.-1 oz., Q. Salieylas T-4 dr. Actions. Ext. Antiseptic-1-1000-250, Int. Stom-Arrests movement of white blood-cells. achic. may, dose ic ac. 4.5%

Increases stability of O—hemoglobin. Lowers temperature. Diminishes metabolism. Small doses stimulate heart. Small doses stimulate brain. Antipyretic, acting on heat centres. Diminishes reflex excitability of cord. Diminishes uric acid and urea. Stimulates uterus. Specific for malaria.

Uses. Ext. Injections—1-4000-500—in amedic dysentery. Int. Stomachic. Tonic. Malarial paroxysms and other malarial manifestations, e.g., neuralgia. Antipyretic. In inflammations gen-

cautions. Do not give in: 1. Middle-ear disease.
2. Gastro-enteritis. 3. To persons with idiosyncrasy for.

CINNAMOMUM SAIGONICUM:

C. Zeylanicum.

Cinnamomum Cassia

Spr. c..... 10-30 M.

COCA (erythroxylon). Paru + Bolivia

Cocainse hydrochloridans 1-4 gr.

Oleature excainse 5 %

came poisoning has resulted for applications of a 2 of a soin to the

her applied excelly it acts on secrel walls on the walls on the contract of the taken into experience of the contract of the c

Actions. Ext. Local anæsthetic. Int. Local anæsthetic, stomach. Antemetic. Vagus depressed. Cerebral stimulant. Respiratory centre stimulated, then depressed. Mydriatic. Accommodation partially paralyzed.

Uses. As local anæsthetic. Gastric pain, vomiting, seasickness.

COCCUS (cochineal):

Use. Coloring agent.

COLCHICUM:

0 11	
C. radix	2-8 gr.
Ext. c. r	1-2 gr.
Px0-6-0%	XX 11.
Windows X	KM.
C. semen	1-5 gr.
Ext. c. s. fl	1-5 M.
Tr. c. s	10-30 М.
Vinum c. s	10-30 11.
Actions. Ext. Irritant, Int. Specific	DE 728
Gastro-intestinal irritant—prostration.	hotagos
Use. Gout and goutiness, including eczen	
ache, bronchitis, etc.	-
aria	
COLOCYNTHIS (bitter calumba)	2-8 gr.
Ext. c	1-2 gr.
* Ext. c. co	5-15 gr.
Pil, cath, co	1-3 pills.
Pil, cath. veg	I-5 "

Actions. Stomachic. Drastric, hydragogue cathartic. Gripes.
Uses. Not given alone. Hydragogue cathartic.
CONIUM (hemlock) Year M. A. 2-5 gr.
Ext. c. fl
Actions. Paralyzes vagus. Ascending paralysis—motor nerves. Mydriatic. Ptosis. Asphyxia. Uses. Rarely used. Whooping-cough, tetanus, chorea, asthma, epilepsy.
Ext. c. fl
Actions and uses. Like digitalis—not so powerful.
COPAIBA 1-1 dr. Macsa σ 1 dr. Oleum c 5-15 Π. Resina c 5-15 gr.
Actions. Of volatile oils generally. Disinfectant, expectorant, and diuretic. Uses. Chiefly in gonorrhœa.
CORIANDRUM (coriander) 10-30 gr. Ol. c 2-5 M.
Actions and uses. Of volatile oils.

	CO10 I-10 gr.
	Abstractum c ½-5 gr.
	Tr. с 1-10 щ.
	Ext. c. fl 2-6 m.
	Cotoin 1-2 gr.
	Paracotoin 1-3 gr.
	Uses. Diarrhœa, ordinary and typhoid or tubercular, Night-sweats.
	CREOLINUM 1-5 m.
*	Action and use. Powerful, non-irritating antiseptic.
	Aqua creosoti
	Actions and uses. Gastric and intestinal antiseptic. Pulmonary tuberculosis (and the broath, much
	CUBEBA (cubeb) 30-60 gr.
	Ext. c. fl
	Oleoresina c 5-30 π.
	** XXX ** X & & X
	Troch. c. (āā 🖁 🎵. oleores.) 1-6
	Oleum c 5-20 π.
	Actions. Of volatile oils. Disinfecting expectorant and diuretic.
	Uses. In cigarettes for bronchitis, asthma, etc.
24	Chronic cystitis, gonorrhœa, etc.
	combine creasote with other drug
Ca	ubante (92 of Cressote) Dose 16. 60m.
4	a some glass of about of the most
-	it than create.

CUPRUM (copper):
C. sulphas (bluestone)dose astringent, \(\frac{1}{2} \) \delta gr. emetic2-20 gr.
Actions. Ext. Astringent. Caustic. Int. Astringent. Direct emetic. Uses. Ext. Caustic and general astringent. Int. Emetic. Diarrhœa. Phosphorus-poisoning.
CURARE (woorari, urari)
Action. Paralyzes motor plates. Use. Tetanus.
CUSSO (kousso, brayera)
DATIIANA ½-1 oz.
Action and use. Aphrodisiac.
DIGITALIS (foxglove). Europe 1-3 gr. 3 Ext. d. 1-1 gr. Ext. d. fl. 1-3 m. Inf. d. Crest consolidation 1-4 dr. Tr. d. 5-30 m. Digitalein. Sol. in. H20. subcut. 100 gr. Digitalin. Proof. in. H20 Gas. disturb. Gas. disturb. Pulse 60 Alabama Alabama Proof.
For high arterest the

to smould lie down after to italis because there is do en stim. the heart & caus SYLLABUS OF MATERIA MEDICA. 40 Actions. Gastro-intestinal irritant. Prolongs diastole, slows rate, increases force, regulates heart. Action direct. Contracts blood-vessels. Action direct and indirect. Raises blood-pressure. Diuretic. (grait kidney allo) Uses. Mitral regurgitation. Mitral obstruction. Aortic obstruction. Tricuspid disease. Cardiac dilatation. In and after acute diseases. Palpitation. Exophthalmic goitre. Hemorrhage. Chronic alcoholism. Acute and chronic nephritis. Caution. Do not use in disease of cardiac musclealls are damaged. DIURETIN.... 15-30 gr. Action and use. Diuretic. DUBOISINÆ SULPHAS (alkaloid from duboisia myoporoides): Action. Like atropine. Use. Mydriatic- 1000 grain disc. ELASTICA (india-rubber, caoutchouc): Uses. Plasters, pessaries, syringes, etc.

ELATERINUM (Squirting Cremital) 10 15 gr.
Trituratio e. 1-t gr.

Actions. The most powerful hydragogue. Gripes.

Much prostration.

Uses. As powerful hydragogue cathartic.

sh. Ergeral miner
THE ACTIONS AND USES OF DRUGS. 41
ERGOTA 30-60 gr.
Ext. e. fl 1-1 dr.
Ext. e 2 . Engotime 5-15 gr.
Vinum e 15 %
Actions. Cardiac depressant. Vaso-constrictor. Hemostatic. Ecbolic. Secretions diminished.
Uses. Cautiously in labor. Post-partum hemor- rhage. Hemoptysis. Hematemesis. Antigalac-
Pois oring (Grangeme (St. authorize fine). ETHYL CHLORIDE.
Use. Spray for local anæsthesia, freezing. Anodyne.
EUCALYPTUS ½-2 dr.
Ext. e. fl
Oleum e 5-30 M.
Eucalyptol 5-30 Ψ.
y 3 x Mun
Actions. Ext. Antiseptic, disinfectant. Int. Stomachic. Stimulates heart. Antipyretic and antiperiodic. Disinfecting expectorant and diuretic. Diaphoretic.
Uses. Ext. As antiseptic. Int. Diphtheria. Fetid bronchitis, cystitis, pyelitis.
EUONYMUS (wahoo, spindle tree) 1-2 dr.

42 STEPADOS OF MATERIA MEDICA.
EUROPHEN. 28% I
Action and use, Antiseptic.
EXALGINUM 1-3 gr.
Action and use. Analgesic.
FABIANA (pichi): [daluable (Dr. S.)] Ext. f. fl
Use. Inflammations of gu. tract. good for stranger
FEL BOVIS (fel tauri, ox-gall) 5-15 gr. F. b. purificatum 5-15 gr.
Actions and uses. Cholagogue purgative. Intestinal antiseptic (?). Enema.
FERRUM (iron):
Ferrum reductum 1-5 gr.
F. sulphas, ½-2 gr.
F. sulphas exsiccatus ½-2 gr.
F. " granulatus \frac{1}{2} gr. K. Wistura f. co. (Griffith's mixture). 2-4 dr.
Pilulæ f. 63 (Blaud's pills) 1-2 pills.
F. carbonas saccharatus 2-10 gr.
Massa f. carbonatis (Vallet's mass) 1-5 gr.
K. iodidum saccharatum 5-15-gr-
Syrupus f. iodidi 5-30 M.
Pilulæ f. iodidi 1-2 pills.
· (Sulphate)
entine of the Suitable fare valuable
monal's astringents

THE ACTIONS AND USES OF DRUGS. 43	
(F. laotas	
F. chloridum.	
Liquor f. chloridi 2-10 M,	
Tr, f. chloridi 5-60 M.	
Liq. f. et ammonii acetatis (Bash-	
am's mixture) 1-4 dr.	
Liquor f. tersulphatis.	
(-" f. nitratis.)2-10 11h	
" f. subsulphatis (Monsel's solu-	
tion) 2-10 M.	
(" f. citratis)	
f. acetatis	
F. exidum hydroxidum 1-4 dr.	
Trochisci 1 1 2	
F. enidum hydrogian magnesiao ido 1-4 dr. = 000.	Lat.
Ferri et ammonii sulphas 2-10 gr.	
Scale preparations:	
F. phosphas solubilis I-5 gr.	
Syrupus f., quininæ et strychninæ	
Phosphatum. dr. 5-15 gr.	
" " ammonii citras 1-5 gr.	
Vinum f. Krikijk, cettradis 1-2 dr.	
F. et strychninæ citras 1-3 gr.	
" " ammonii tartras 5-15 gr.	
F. citras 1-5 gr.	
n 13 Jenie phosphate 190 3 3 3	
of your	
Strychning & "	
32	

FERRUM DIALYSATUM..... 10-30 M.

Use. Arsenic-poisoning.

FICUS (fig)..... freely,

Use, Mild purgative.

FORMALDEHYDE (formalin):

Commercial preparation is a solution of the gas.

Uses. Ext. Antiseptic and disinfectant for inanimate. objects. Hardening agent ils histological tech-

1 6 % formalin per 1 to auft.

FRANGULA (buckthorn)..... +- t dr. Ext. f. fl..... +2 dr.

Action and use. Mild purgative.

get mid of odor spray with 11 #484 fame so formed composed in the w

1/2 gr. in \$1.0 on an empty al

flandulae uprarenales siresa	492
hyroid is - powerful vaso dilator.	7 gr.
THE ACTIONS AND USES OF DRUGS. 45	
GALLA (nutgall), tannic acid 50-60%.	
Tr. g 1-2 dr.	
Ung. g	
Actions and uses. See tannic acid.	
GELSEMIUM (yellow jasmine) 5-10 gr. Ext. g. fl 5-10 ml.	
Tr. g ‡-1 dr.	
Gelsemina 1 1 20 gr.	
Actions. Paralysis from depression of anterior cor- nua. Depression of sensory tracts (?). Mydriatic.	
Diplopia. Asphyxia. Uses. Neuralgia, migraine. Mydriatic.	
GENTIANA (gentian) 5-30 gr.	
Ext. g	
Tr. g. co	
Action and use. Stomachic bitter.	
GLONOIN. See Spirit of.	
GLYCERINUM 5-60 П.	
Glyceritum amyli serecutuit 31 % Bones a	
Suppositoria glycerini. p. r. n.	
Actions Firt Solvent vehicle Int. Purges De-	
mulcent. Sweeting agent in win	hales 11
1 Gambage	5 gr.
ilulas catharticas Co.	5-1
tions - Drestie hydrago que colhe	

Uses. Ext. Vehicle. Chapped hands. GLYCYRRHIZA (licorice root): Ext. g..... freely. Ext. g. purum..... Ext. g. fl 1 dr. Glycyrrhizinum ammoniatum 5-15 gr. Pulv, g, co..... 1-2 dr. Mist, g. co. (Brown mixture)..... 1-4 dr. es. Demulcent, vehicle. Pulv. g. co. as laxative. Mist, g. co. as expectorant. GOSSYPII RADICIS CORTEX (cotton root bark..... 1-1 dr. Actions and uses. Like ergot. GOSSYPIUM PURIFICATUM (absorbent cotton). GRANATUM (pomegranate) 1-1- dr. Actions and uses, Astringent, Anthelmintic for Felletterne tamore official Pulleterint RINDELIA 1-1 dr. Ext. g. fl 1-I dr. Actions. Expectorant, Relaxes muscle of bronchial tubes.

Uses.	Ext.	Poison	ivy.	Int.	Asthma,	chronic bron-
chit	tis wit	h asthm	atic	sympt	oms.	

of, GUAIACOLUM 2-10 M.
G. benzoas 2-10 gr.
G. carbonas 5-30 gr.
G. salicylas 5–30 Π.
Actions and uses. Like creosote. Less of a gastro intestinal irritant.
GUAIACUM:
C lignum (lignum vita) 1-1 dr.
G. resina (guaiac). Quaiacum 5-30 gr.
Tr. g 1-1 dr.
Tr. g. ammoniata ½-1 dr.
Actions. Stomachic, increases peristalsis. Purgative.
Uses. Mild purgative. Empirically for sore throat
Immilitie - Sozenge of quaine + quare gelly.
GUARANA (Brazilian cocoa) 1-1 dr.
Ext. g. fl ½-1 dr.
Action. Like caffeine. Use. Chiefly sick headache.

GUTTA PERCHA.

Liq. gutta percha.

Uses. Making splints:

HÆMATOXYLON (logrocod): Ext. h 5-15 gr.
Action and use. Ext. and int. Astringent.
Hamameligit Tolia
HAMAMELIA (vitchhazel) Contact 30 and HExt. h. (bliorum 1-2 dr. agent Hamamelisis Action and use. Ext. and int. Astringent.
Action and use. Ext. and inc. Astrugent.
HEDEOMA (pennyroyal); Oleum h
Actions and uses. Of volatile oils,
Actions and about of tomore only
HIRUDO (kech):
Use. To draw blood.
HOMATROPINÆ HYDROBROMALUM
(1) H +0) Tiso =0 subcutaneously.
110 20
Actions and uses. Like atropine.
HUMULUS (hops):
DE DANIEL X-X dr.
Lupulin 5-30 gr.
Ext. 1. fl 5-30 M.
Oleores, 1
Actions. Stomachic, tonic, carminative, soporifie, narcotic,
Uses. Dyspepsia, insomnia, delirium tremens.

HYDRARGYRUM (mercury):

Actions in common of all preparations. Ext.

Stimulating, irritant to caustic. Nearly all, antiparasitic and antiseptic. Relieve itching. Absorbed by skin; by lungs in form of vapor. Int.
Gastro-intestinal irritants in varying degrees.
Converted into complex albuminate in stomach,
becoming an oxyalbuminate in blood. Large
doses produce anemia. Indirect cholagogues.
Excreted in saliva, bile, urine, milk, sweat. May
salivate. Specific for syphilis.

FOR EXTERNAL USE ONLY.

Empl. ammoniaci cum h.

Use. Local application.

Ung. h... Blue on the 50% 31 4 of a Actions. Antiparasitic. When absorbed, like other mercurials.

Uses. Antiparasitic. Application to syphilitic and other sores. Inunction for chronic joint and other inflammations; enlarged lymph nodes; to rapidly mercurialize.

H. ammoniatum (white precipitate).
Ung. h. ammon. (white precipitate ointment).

Uses, Antiparasitic. Syphilitic and other sores.

H. oxidum flavum.

Ung. oxide flavin

20 gime by huppo.

Man 10 % and and Hy Saliculate in all the first of the control of the saliculate in all the control of the

en which is behaveled by the discharge

50 SYLLABUS OF MATERIA MEDICA.

Oleatum h.

Uses. Antiparasitie. Inunction.

Ung. h. oxidi rubri (red precipitate sintment).

Uses, Antiparasitic. Syphilitic and other sores.

Liq. h. nitratis.

Uses. Caustic for syphilitic warts, condylomata, etc.

Ung. h. nitratis (citrine cintment).

Uses. Caustic. Diluted, antiparasitic. Syphilitic and other sores.

FOR INTERNAL AND EXTERNAL USE.

H. cum creta (mercury with chalk), 38%.. \(\frac{1}{2}\)-10 gr.

Massa h. (blue mass, blue pill), 33%.....\(\frac{1}{2}\)-15 gr.

Actions. Int. Laxatives or purgatives. Indirect cholagogues.

Uses. Chiefly as laxatives or purgatives or cholagogues. Mercury with chalk for vomiting, hepatic congestion, cirrhosis, diarrhoea, Blue mass in diuretic pill or for syphilis.

Lotio nigra (black touth), not offi-

co est colorynth -- 15 "

Actions. Ext. Mildly irritant and stimulant. Absorbed by skin, by lungs. Int. Laxative and purgative-duodenum and jejunum-spinach-green or dark stools. Gripes. Indirect cholagoguepartly converted to bichloride, hence also direct.

Uses. Ext. Stimulant for syphilitic and non-syphilitic sores. Itching. Vapor bath in syphilis. Int. As general laxative or purgative-at night, saline in morning. Vomiting from congestion of stomach. Congestion of liver, functional or organic. Cirrhosis. Diarrhœa. Diuretic in small doses or in "diuretic" pill (calomel, squill, digitalis, āā gr. 1). Rarely for syphilis.

* H. chloridum corrosivum (corrosive chloride, bichloride) gr. Lotio flava (vellow zwash), not official, 2 gr. to I oz.

Actions. Ext. Antiseptic, disinfectant, antiparasitic -1-70000-500. Absorbed by skin, by lungs. Int. Gastro-intestinal irritant. Produces anemia

in large doses. Direct cholagogue.

Uses. Ext. As general antiseptic and parasiticide. Yellow wash for itching and syphilitic sores. Int. Alone or with potassium iodide for syphilis-2d or 3d stage. To absorb non-syphilitic, chronic, inflammatory formations-e.g., interstitial nephritis.

H. iodidum flavum (yellow, green, or protiodide) 1-I gr. H. iodidum rubrum (red or biniodide) .. 10 10 gr. Lig. arseni et h. i. (Donovan's solution), 1% of each..... I-10 M.

H. oxidum robrum (red precipitate).... 1-1 gt

Uses. Chiefly internally for syphilis. Donovan's so

-H. subsulphas flavus (turpeth mineral)... 2-4 gr Use. As emetic

MODES OF ADMINISTRATION.

- I. Inunction-blue ointment and oleate chiefly,
- 2. Endermically-calomel.
- 3. Hypodermically-bichloride.
- 4. Fumigation-calomel chiefly.
- 5. Suppositories-blue ointment chiefly.
- 6. Baths-rarely-bichloride in water.
- 7. Inhalation-rarely.
- 8. Internally-most usual-any preparation for in

*HYDRASTIS (golden seal)	5-60 gr
Ext. h. fl	5-60 m
Tr. h	1-2 dr
Glyceritum hydrastis	5-60 M
Hydrastininæ hydrochloras,	1 gr
Berberine.	をかった

Actions. Stomachic. Cholagogue, Diuretic. Stim ulates involuntary muscle. Antiperiodic.

Dest aubstitute for ergot

Uses. Ext. Lotion in chronic inflammations. Injection in leucorrhœa, etc. Int. Chronic inflammations of mucous membranes. Menorrhagia, metrorrhagia, Malaria, Wering HYOSCYAMUS (henbane). Ext. h Ext. h. fl.... Hyoscyaminæ sulphe... hydrobromac um Hyoscinæ Actions. Like belladonna and stramonium. Differences: 1. Cerebral and spinal sedative. 2. Relieves griping better. 3. Relaxes muscle of urinary tract more. Uses. Chiefly with other purgatives and for vesical and urethral spasm, cystitis, etc. Hyoscine for acute mania, delirium tremens. Chorea, paralysis agitans, etc. Hyoscyamine for sea-sickness, author HYPNAL (chloral and antipyrin)...... 15 gr. Actions and uses. Hypnotic and analgesic. HYPOPHOSPHITES: Calcii hypophosphis..... Sodii Potassii Syr. hypophosphitum....

and the second s
Zinci phosphidum in 1 gr.
Acidem hypophosphorosum dilutum. 10 %
Actions and uses, Chronic bronchitis, Pulmonary
tuberculosis. Reconstructive.
1CHTHYOCOLLA (Isinglass) freely.
Empl. i. (court plaster).
ON CONTRACTOR OF THE PARTY OF T
Uses. Emollient and protective.
WHTHVOI - amminut intelliged sulphoniste
Uses. Ext. Chronic eczema and psoriasis Int.
Usas But Cherrie and antiseptic
Chronic rheumatism. Inter cuttachtic
Silver in the same of the same
IODOCODMUM 9 5to .17
10D0F0RMUM9.539 1-3 gr.
Ung. iodoformi 10%.
Actions and uses. Ext. Antiseptic, disinfectant,
analgesic. Int. Unknown.
= setratodopyrral
HODOLUM 787, I. 9070 1-3 gr.
A STATE OF THE PARTY OF THE PAR
Action and use. Like iodoform. Sometimes used
internally.
IDDUM: (Uttained from askes of seame
Liq. iodi co. (Lugol's solution) I 5%, KI 10%1-10 M.
Tr. iodi, 7% 1-5 M.
Ung. iodi, 4%.
smilk for I korening.
emularon of starch + milk
nancos

ucac +	has a special action on dysentery.
emayo	give 10 m. landamin to he before
in after	sec . O forment wounder
56	SYLLABUS OF MATERIA MEDICA.
V _A	ctions. Ext. Irritant. Antiseptic. Int. Stomachic, emetic (direct and indirect). Expectorant. Cholagogue. Diaphoretic.
1	ses. Ext. Anthrax. Int. Stomachic. Vomiting of pregnancy. Gastric ulcer. Slow emetic in bronchitis, etc., in children. Dysentery. Hepatic dyspepsia Bronchitis, astama. Diaphoretic.
15	RIDIN (irisin, from blue flag) 1-5 gr.
A	ction and use. Cholagogue.
11	(15 (blue flag) 10 30 gr.
	Ext. 1
A	ction and use. Cholagogue purgative.
* J	ALAPA Merico 5-20 gr.
bitarhate	ALAPA Merico 5-20 gr. Fat. j. 2-8 gr. Pulv. j. co. = Pulvio Paragano 1-1 dr. Res. j. 1-5 gr.
	ses. Nephritis, uræmia, dropsy etc. cardine
JI	EQUIRITY.
7	ction Ext. Intense local inflammation, with con- stitutional reaction. ses. Granular lids. Certain skin diseases. Lupus, Epithelioma, etc.
posit o	five Jalap and Hopel in a filt. Wal

ne or two powders. Agel may bean Hylls

Aqua danocerasi = Cherydond water Strength = 0.4% whoolute HCM.

THE ACTIONS AND USES OF DRUGS,	57
JUGLANS (butternia)	dr.
Ext. j 5-30	gr.
Action and use. Mild cathartic. Resembles a	rtiu-
Kaolin (al. siliente)	
KAFIALA (rottlera) 1-2	dr.
Actions and uses. Anthelmintic for tapeword threadworm, roundworm. Purgative.	rm,
KRAMERIA (rhatany) 5-30	gr.
Ext. k 5-10	
Ext. k.fl 5-30	TIL.
Tr. k 1-2	dr.
Troch. k. (1 gr.)	
Syr. k 1-4	dr.
Actions and use. Ext. and int. Astringent, throat, epistaxis, gonorrhoea, diarrhoea, etc.	Sore
LACTUCARIUM (lettuce) 5-60	gr.
Tr. 1 1-2	dr.
Syr. 1	dr.
Action and use. Hypnotic. Mydriatic.	
LIMONIS CORTEX (lemon peel):	
SOND	dr.
Oleum I.	
Jr. himmin Conticio	
Leptandra Jahreis Rott) 4 - 1 3	
The sale.	an

Uses. Vehicle.	Flavoring agent.
----------------	------------------

The state of the s
LIMONIS SUCCUS (lemon juice) 1-2 oz.
Action. Like citric acid. Uses. Thirst. Effervescing mixtures. Scurvy.
LINUM (linseed):
Oleum 1 1-2 oz.
Uses. Poultice. Burns. Demulcent.
LITHIUM. C. carbonas. L. citras. L. citras effervescens. L. citras effervescens. L. citras and uses. Solvents of uric acid, diuretics, and alkalinizers of the urine in gout and allied conditions. N.A.
No.

Actions. Gastro-intestinal irritant, with intense prostration. Relaxes muscle of bronchial tubes. Uses. Asthma, and bronchitis with asthmatic symptoms.

LYCOPODIUM.

Uses. Dusting powder, etc.

LYSOLUM.

Action and use. Antiseptic.

Action and use. Like volatile oils.

MAGNESIUM:

M. sulphas (Epsom salt)	1-1	oz.
M. carbonas		
M.Schros effervescens	1-1	oz.
Liquor magnesii citratis		
Magnesia ponderosa	5-60	gr.
Magnesia oxidim	5-60	gr.

Actions. Magnesia and magnesium carbonate, antacid. Sulphate, carbonate, and citrate, saline purgatives. Diuretic; alkalinize blood and urine (feeble).

Uses. As alkalies and antacid. Antidote to mineral acids, oxalic acid, salts of mercury, arsenic, copper, and alkaloids. Sulphate antidote to lead and barium. Purgatives, Gout and gravel.

MALTUM:

Ext, m..... 1-4 dr.

Action and use. Food.

MANGANUM:
M. dioxidum. Praci pitatom 2-10 gr.
M. sulphas 2-8 gr.
Potassii permanganas 1-2 gr.
Actions. Ext. Disinfectant, deodorant, antiseptic. Int. Not known. Uses. Ext. As above. Salivation, gonorrhea, leucorrhea, uterine disease, etc. Int. As hematinic. Amenorrhea
newly is an contido to for morphime.
MANNA 1-2 oz.
Use. Laxative.
MASTICHE (mastic):
Pil. aloes et m. ("Lady Webster") 1-3 pills.
Uses. Mastic, for varnishes, etc. Pil. aloes et m. as laxative.
MATRICARIA (Ger. chamomile) 1-1 dr.
Action and use. Stomachic bitter.
MEL (honey): M. despumatum depenature freely.
Uses. Demulcent. In cough mixtures. Laxative in children.
MELISSA (balm) 1-2 ds.
Use. Flavoring agent.

the state of the same of the same

THE ACTIONS AND USES OF DRUGS, OI
MENTHA PIPERITA (peppermint):
Spr. m. p 5-15 M.
Ol. m. p
Aqua m. p ½-2 oz.
7x0xx.xi.xg.x, xexx
Actions. Ext. Of volatile oils. Anodyne. Int. Stomachic, carminative. Uses. Ext. Neuralgia. Int. Carminative, stomachic, flavoring agent.
MENTHA VIRIDIS (spearmint):
Spr. m. v 5-15 η.
Ol. m. v 1-5 M.
Aqua m. v 1-2 oz.
Actions and uses Of peppermint.
MENTHOL (from peppermint) ½-2 gr.
Actions. Antiseptic, local anæsthetic, anodyne. Uses. Spray in common cold. Neuralgia. Asthma.
METHYL CHI ODIDE
METHYL CHLORIDE.
Use. Spray for local anæsthesia, freezing. Anodyne.
METHYLENE BLUE - mathethioninae methylene Blue - mathethiodiny - 5 gr.
Action and uses. Rheumatism, myalgia. Vesical irritation.
n and state of the stranger
the france
prevent strangung me methyle

Jo

strychine from Strychnos ignation

	1 12 perfermes advances
	62 SYLLABUS OF MATERIA MEDICA.
5	METHYLENUM DICHLORIDUM.
	Action and use. General anæsthetic.
	METHYLLS ALICYLAS (artificial oil of winter- green)
- 6	Actions and uses. Like salicylic acid
	Tr. m
	Action. Diffusible stimulant. Uses. Collapse, fainting, etc. Typhoid, pneumonia. Hysteria,
	MUSCARINA (alkaloid from amanila muscaria, fly fungus) 1/8-2 gr.
	Action. Like Calabar bean. Decreases force and frequency of heart. Lengthens diastole. Produces dyspnœa, paralysis. Sialagogue. Diaphoretic. Stimulates involuntary muscle.
	Uses. Rarely used. To increase peristalsis. In- flammatory exudates.
3	MYRISTICA (nutmeg)
un of V	Action and use. Like volatile oils.
- int	ition that has carred angine per
eard	ation that has caused argume per

63 THE ACTIONS AND USES OF DRUGS. MYRRHA dr. Tr. aloes et m...... 1-24 Pil. aloes et m..... 2-10 pills. Actions and uses. Of volatile oils. Expectorant, uterine stimulant, emmenagogue. NAPHTALENUM 1-20 gr. Action. Local. Uses. Intestinal ulcerations-dysentery, typhoid, tubercular. NAPHTOL (beta-naphtol)..... 1-20 gr. Action and use. External and internal antiseptic. NITROGLYCERINE. See Spirit of glonoin. Doce must be as large as 10- 1, gr (2-5 m) NUX VOMICA (poison nut, dogbutton). . 1-4 gr. Ext. n. v. fl...... 1-4 Strychnina.... (. 4.700 . + 70.). 1 1 gr. Actions. Stomachic, increases peristalsis cardiac stimulant. Blood-pressure raised. Stimulates respiratory and vaso-motor centres. Spinal convulsions. Sharpens sight and hearing. Uses. As stomachic and carminative. Constipation. General cardiac stimulant. Respiratory stimulant. mica is a better &

64 SYLLABUS OF MATERIA MEDICA.
OLEUM BETULÆ WORATOR 1-5 T
Actions and uses. Like salicylic acid.
OLEUM CADINUM:
Action and use. Like tar. Skin diseases, e.g. psoriasis, chronic eczema, pruritus.
OLEUM GAULTHERIÆ (oil of wintergreen), 1-5 m Spr. g
Actions and uses. Like salicylic acid.
OLEUM GOSSYPII SEMINIS (cotton-seed oil).
Uses. As other fixed or neutral oils.
OLEUM JUNIPERI. 2-10 M Spr. j. 2-10 M Spr. j. 1-1 dr Spr. j. 1-4 dr
Actions. Like turpentine. Use. Diuretic in heart disease, ascites, nephritis.
OLEUM LAVANDULÆ FLORUM (oil of laven

Actions and uses. Stomachic, carminative, flavoring agent.

OLEUM MORRHUÆ (cod-liver oil) 1-4 dr. Action. Ext. Bland oil. Int. Easily absorbed food. Uses. Pulmonary tuberculosis. Rickets. Chronic bronchitis. Chronic wasting diseases. Tertiary syphilis, etc. Contra-indications in pulmonary tuberculosis: 1. High temperature. 2. Severe hemoptysis. 3. Dyspepsia. 4. Vomiting. 5. Diarrhea.
OLEUM OLIVÆ (olive oil) freely. Actions. Ext. Protective. In massage. Int. Pro-
tective. Sedative. Laxative. Uses. Ext. Burns. Int. Corrosive poisoning. Gall stones. In enemata. **Corrosive poisoning. Gall stones. In enemata. Hypercalls that a stone of the stone of
Action and uses. Like turpentine. Pleasanter. OLEUM RICINI (castor oil)
Actions. Ext. Protective. Galactagogue. Int. Simple laxative or mild purgative—5 hrs. Uses. Universal laxative or mild purgative. Galactagogue. tagogue. Stumbalis prequant nterms.
OLEUM RUTÆ (oil of rue) 1-4 M. Actions. Ext. Irritant, vesicant. Int. Gastro-in-
testinal, renal, ovarian, and uterine irritant. Uses. Rarely given. Que no en la accompant. Jos en parcheo hydria turnishes mutrimunt.
Junishes mitimut

	66 SYLLABUS OF MATERIA MEDIC	Α.
	OLEUM SANTALI (sandal-wood)	2-10 П.
	Actions. Of volatile oils. Stimulating expectorant, and diuretic.	disinfectant,
	Uses. Gonorrhœa, etc. Chronic bronch	itis.
	OLEUM TEREBINTHINÆ (oil of tur	rpentine),
	dose { anthelmi	5-30 M. ntic, ½-4 dr.
	Ol. t. rectdose anthelmi	5-30 M. ntic, ½-4 dr.
	Actions. Ext. Rubefacient, irritant, courselver, Disinfectant. Int. Storminative; purgative in large doses. To tapeworm; to threadworm as energy stimulant, directly, and reflexly fro Vaso-constrictor. Raises blood-prepresses nervous system in large dose expectorant. Stimulates gu. mucous Hematuria. Urine smells of violets rus-poisoning. The Anthelmintic. intestinal hemorrhage. Hemoptysis. gangrene. Chronic bronchitis.	nachic, car- Anthelmintic na. Cardiac m stomach. ssure. De- s. Powerful s membrane. s. Phospho- Neuralgia, Gastric and Pulmonary
	OLEUM THEOBROMATIS (cacao butt	
	Use. Making suppositories.	.,
	OLEUM THYMI	I-5 M.
	Actions. Like copaiba.	
- 9	wen .	Da
Z. Ca	Rente of anger	10 .
5. By	R	
. ou	. I must of angor	

Uses. Bronchitis. Gonorrhœa, leucorrhœa, etc
gudia & Phillipines
OLEUM TIGLII (croton oil) 1-2η.
Actions. Ext. Powerful irritant. Pustulation. Int. Powerful purgative—I-2 hrs. Produces enteritis.
Uses. Ext. Rarely used now. Int. Purgative in obstinate constipation, fæcal impaction, etc. Apoplexy.
Cautions. Do not give to: 1. Children. 2. Pregnant women. 3. Feeble persons. 4. In hemorrhoids. 5. Peritonitis. 6. Gastritis. 7. Enteritis. 8.
itis. 8. Organiza abstruction.
OPIUM (from poppy) 1-2 gr. Alkaloids:
Morphine, codeine, thebaine, narcotine, nar- ceine, and 14 others.
Opii pulvis 1-2 gr.
Ext, o
Troch, glycyrrhizæ et o 1-4
Vinum o. (Sydenham's laudanum),
10% 3-20 M.
Pil. o., āā 1 gr 1-2 pills.
Pulv. o. et ipecacuanhæ (Dover's
powder), 10% 3-15 gr.
Acetum o. (black drop), 10% 3-20 Ψ.
Opium granulatum - 1 gr.

SYLLABUS OF MATERIA MEDICA. Tr. o. (laudanum), 10%..... 3-20 III. Tr. o. camphorata (paregoric), 0.4%, 1-4 dr. gr. Tr. o. dead, 10%..... 3-20 M. Tr. ipecac. et o..... 3-15 m. Morphina. 4. 3. 5.0. 42.0 . . . 1 OT. M. hydrochloras 24 1 M. acetas 2:67 ... 1-1 M. sulphas..... 15. 2...... +1 Pulv. m. co. (Tully's powder), I gr. to I dr..... 5-15 gr. Trock, m. et ipecac., an gr.... 1-5 Magendie's solution: morphia sulphate, 16 gr.; water, I oz. U. S. Ph. solution: morphia sulphate, I gr.; water, I oz. ions. Ext. Local anodyne. Absorbed by oraw surfaces. Int. Stimulates heart. Slows pulse. First excites, then calms psychic centres. Narcotic, Produces sleep. Abolishes pain. Depresses respiratory centre, slowing respiration. Depresses anterior cornua. Decreases all secretions except sweat. Diaphoretic. Antispasmodic. Myotic-"pinhole pupil." Diminishes sugar in some cases of diabetes. > peristalsis Uses. Ext. Anodyne, Int. General anodyne, Gastric or intestinal pain. Diarrhœa, Cardiac or aneurismal pain. Insomnia in selected cases. In cough mixtures. Diaphoretic in common cold.

doses produce no THE ACTIONS AND USES OF DRUGS. Ornet of severe acute infections Antispasmodic in renal and biliary colic. To stop peristalsis in peritonitis, etc. Diabetes. W.3. V. Cautions. I. Beware of opium habit. 2. Give cautiously to children, and, 3, in kidney disease. > hight aments of phthieses CODEINA (80 420) 1-2 gr. Actions." Like morphine. Excites cord, producing tremors. Depresses psychic centres less than morphine. Uses. Analgesic. Cough mixtures. Diabetes. OUABAIN...... foo gr. Action. Local anæsthetic. Use. Like cocaine. PANCREATINUM..... 5-15 gr. Uses. Artificial digestive agent. PARALDEHYDUM 1-1 dr. Action. Powerful hypnotic. . Uses. Hypnotic. Mania, melancholia, PENTALUM. Action and use. General anæsthetic. PEPO (pumpkin seed)..... 1-3 oz. Action and use. Anthelmintic for tapeworm.

Personal and the Control of the Cont
PEPSINUM 5-60 gr.
-P. saecharatum 5-60 gr.
Uses. Digestive agent in or outside of body.
PETROLATUM LIQUIDUM:
" mod (soft).
Actions and uses. Protective and vehicle. Spray.
walphenacetinum 5-10 gr.
Actions. Antipyretic, analgesic. Cardiac depressant.
Uses. Reducing temperature. Neuralgic and other
present than autamilia + antipyrin
PHOSPHORUS 1 100 20 gr.
Oleum phosphoratum, 1% 1-5 呗.
→ Pilulæ p., ãã 100 gr
Spr. p 8-40 m.
Elixir p ½-2½ dr.
Actions. Int. Increases strength of bones. Uses. Int. Osteomalacia. Rickets. Ununited fracture. Alterative. Habitat tropical W. Ofrica
PHYSOSTIGMA (Calabar bean) \(\frac{1}{2} - 2 \) gr. Ext. p \(\frac{1}{10} - \frac{1}{2} \) gr.
Tr. p
P. sulphas (eserine s.). New 100 - 10 gr.
Loo gr
1 90
3/ "
- /

Actions. Rate of heart slowed, force increased. Blood-pressure raised. Depresses anterior cornua, Sialagogue, hidrotic. Stimulates involuntary muscle. Myotic, spasm of accommodation, decreases intraocular tension. Uses. Tetanus. Strychnia-poisoning. Chronic constipation. Vesical atony. Chronic bronchitis. As myotic. Glaucoma. Paralysis of iris and ciliary muscles.
PHYTOLACCAE FRUCTUS (pokeberry) 5-30 gr.
Phytolacca RADIXdose { emetic 5-30 gr. alterative, 1-5 gr.
Ext. p. radicis fl 5-30 π.
Actions and uses. Emetic and purgative.
PICROTOXINUM 120 40 gt.
Actions. Ext. Parasiticide. Int. Unknown. Poi-
Uses. Ext. Pediculi. Int. Sweats of pulmonary tuberculosis.
PILOCARPUS (jaborandi). Brogil 5-60 gr.
Ext. p. fl 5-60 m.
Pilocarpine hydrochloradum 1 g gr.
Actions. Cardiac depressant. Stimulates all secre- tions except bile. Powerful diaphoretic. Stimu- lates involuntary muscle. Myotic.
Uses. With caution. Nephritis: Antidote to bella-donna-poisoning.
est quie bimetics
cade by hugher of strugch or eath
estal injec. of atmospher.

PIMENTA (allspice)
Action and uses. Like caryophyllus.
PIPER (black pepper) 5-20 gr. Oleoresina p 1-1 gr. Piperinum 1-10 gr.
Action and use. Like volatile oils.
PIPERAZINUM
PIX BURGUNDICA: Empl. p. b. Empl. picis contheridatum.
Use. Basis for plasters.
PIX LIQUIDA (tar)in pills ‡-1 dr. Syr. p. l
disinfectant, expectorant. W. and Late Chronic bronchitis, pulmonary tuberculosis.

THE ACTIONS AND USES OF DRUGS.

PLUMBUM (lead):

P. oxidum.

Emplastrum plumbi.

· Unguentum diachylon.

Liquor plumbi subacetatis (Goulard's ext.).

Liquor plumbi subacetatis dilutus (lead water).

Ceratum plumbi subacetatis (Goulard's cerate).

* XXXXXXX

Digital September 1980 Marks 1980 Marks

P. nitras.

P. iodidum.

Uxgrenkunkphinnel.indidiak

Actions. Powerfully astringent. Hemostatic. Both int. and ext.

Uses. Ext. As general astringent. Int. Diarrhœa, gastric and intestinal hemorrhage.

PODOPHYLLUM (may-apple) Wents with 5-20 gr.

Ext. p. fl. 5-20 m.

Res. p. (podophyllin). 1-1 gr.

Use. As cholagogue purgative. - a cholagogue purgative.

Pil, Podophylli, Bel. et Capsici - 1 &

gest of Pad . - 14 gr. 7 (Cape sel. leaves 18 ...

73

POTASSIUM:

Liquor potassa.	hydroxidi	5-30	m freely	diluted.
Potassa/cupis				

Actions. Ext. and int. Irritant, caustic, antacid.
Gastric sedative. Alkaline diuretic. General
muscular depressant.

Uses. Ext. As 'a caustic, int. Gastric sedative.
Alkaline diuretic.

P.	carbonas						*	*		*	+.			i	5-30 g	r.
P.	bicarbonas				*				*						5-60 g	r.

Uses. Gastric sedative. Acid eructations.

P. acetas	5-60 gr.
P. citras	5-30 gr.
Liquor p. citratis	1-8 dr.
P. citras effervescens	5-60 gr.

Actions. Diuretic, alkalinizer of urine, diaphoretic. Uses. Gout, uric acid gravel, scurvy, diuretics generally.

P. sulphas	1-4 dr.
P. bitartras (cream of tartar),	

dose { diuretic and refrigerant, 20-60 gr. purgative..... 1-4 dr.

Actions. Saline purgatives. Sulphate a cholagogue.
Bitartrate an alkaline diuretic and diaphoretic.

Uses. Sluggishness of bowels. Bitartrate as a general diuretic and diaphoretic,

P. nitras (saltpetre, nitre) 5-20 gr., Cherta potassii nitratis.
Actions. Gastric irritant. Cardiac depressant. Di- uretic. Diaphoretic. Uses. Diuretic. Diaphoretic. Asthma.
P. chloras
Actions. Gastric irritant. Disintegrates red corpuscles. Paroxysmal hemoglobinuria. Death, heart failure.
All potassium salts are cardiac depressants. Potassium salts are said to be hematinic. All alkalies, given before meals, increase secretion of hydrochloric acid; given after meals, neutralize some of acid secreted.
PRUNUM freely.
Use. As laxative. East M A PRUNUS VIRGINIANA (wild cherry) ½-1 dr. Ext. p. V. fl. ½-1 dr. Infusum p. V. 1-2 oz. Syr. p. V. 1-4 dr.
Action. Like dilute hydrocyanic acid. Uses. Cough mixtures. Vehicle,

76 SYLLABUS OF MATERIA MEDICA. Action. Like senega. Uses. Asthma. Chronic bronchitis. Orchitis, epididymitis. PYROGALLOL (pyrogallic acid). Actions, Ext. Irritant, Parasiticide, Uses, Ext. Do not apply over extensive surface. Chronic psoriasis. Chronic eczema. Ringworm. PYROXYLINUM (guncotton): Collodium (collodion). .. flexile. blister in 8 th. (blistering collodion). Produc Uses. Protective and solvent. QUASSIA: Ext. q..... 1-3 gr. Tr. q..... 1-2 dr. Quassin..... 1 gr. Actions and uses. Stomachic bitter. Injected, anthelmintic to threadworm.

OUERCUS ALBA (white oak) contains 6-11% tannic acid.

3e ext ..

QUILLAJA (soap bark) 1-1 dr.
Tr. q
Uses. Mixing oily liquids with water. Expectorant.
RESINA:
Ceratum r. (basilicon ointment). Empler. (adhesive plaster).
Action. Antiseptic, stimulant. Uses. Sores, ulcers, wounds, etc.
RESORCINGM 2-5 gr.
Actions and uses. Chronic skin diseases. Sebor- rhœa sicca. Gastric fermentation. Marked car- diac depressant.
West U.S.
RHAMNUS PURSHIANA (cascara sagrada),
Ext. r. p. fl
Action and use. Stomachic and mind purgative.
china
RHEUM (rhubarb)dose { stomachic, 1-5 gr. purgative, 10-30 gr. (4-8
Ext. r 3-%5 gr.
Ext. r. fl 1 dr.
Pil. r., āā 3 gr
Pil. r. co., āā 2 gr
Burparpi distante
Co Printfied also
Coile munt fair

Tr. rdose stomachic, 1-1 dr. purgative, 1-4 dr.
Tr. r. arom 1-3 dr.
CD3 Tr. r. dulcis 1-1 oz.
Mist, r, et sodæ d_2 oz.
9-pacae Syr. r 1-4 dr.
Syr. r. arom dr. for child.
Actions. Stomachic. Purgative—4 to 8 hrs. Gripes. Fæces yellow. Constipation afterward. Urine yellow. Cholagogue. Uses. Stomachic and purgative.
acute fevers.
RHUS TOXICODENDRON (poison 127) 1-5 gr.
Uses. Dangerous. Skin diseases. Nocturnal incon- tinence.
ROSA:
R. gallica (red rose).
Confectio r 30-60 gr.
Ext. r. fl 1-1 dr.
Mel r freely.
Syr. r 1-2 dr.
Richard Aphily A
Oleum rosæ (attar of rose).
A CONTRACTOR OF THE PARTY OF TH
Aqua r
Aqua r. indefinite. Aqua r. Ung. aquæ r. (cold cream). Spermaesti 12.
Uses. Vehicles and flavoring agents. Expres, o'd plumed to the Refortion to
na priete 3

RUBUS (blackberry): \$\frac{1}{2} \text{ dr.}\$ Ext. r. fl	
Use. Diarrhœa.	
RUBY X ID (AXXX (raspberry): indefinite. Use. Flavoring agent.	
RUMEX (yellow dock)	
SABINA (savine)	
SACCHARIN (glusidum) Bergonde 2 gr. Use. Sweetening agent. To replace sugar in diabetes. SACCHARUM (cane sugar)	*

u least wit a Salizin next. 80 SYLLABUS OF MATERIA MEDICA. SACCHARUM LACTIS (sugar of milk), indefinite Uses. Vehicle. Making triturations. Nutrient. SALICINUM 28 Hab (Willed) Acidum salicylicum ... 4. 5.0 . H. D. 5-60 gr. Sodii salicylas 5-60 gr. Actions. Ext. Salicin and salicylic acid, antiseptic. Lanhidrotic. Int. Irritant. Salicin converted to salicylic acid in intestine. Impurities are cardiac derressants. Antipyretic. Antiperiodic. Uses. Ext. Stimulating antiseptic. Removing corns. etc. Excessive sweating. Int. For rheumatic fever. Chronic rheumatism. Migraine. Sciatica. Diabetes. Pleurisy. Cystitis. erachie rolution for washing out canties. SALIPYRIN (antipyrin and salicylic acid). . 6-40 gr. Action and use. Sciatica. Chronic rheumatism. SALOL (phenylis married) (almost med.) 5-30 gr. Actions. Split in intestine into salicylic and carbolic acids. Intestinal antiseptic. Like salicylic acid. Uses. Diarrhœa, dysentery, cholera, chiles. Antiseptic in ureter, bladder, etc. it take salinglated except in form of salot SALOPHEN (almost mode) 5-15 gt. Action. Split in intestine, forming salicylic acid. Uses. Like salicylic acid.

Thierach's solution & Done and 12
. H20 92 me 1000
THE ACTIONS AND USES OF DRUGS. 81
SALVIA (sage) ‡-1 dr.
Uses. Condiment. Night sweats.
SANGUINARIA (bloodroot). W.A. 2-30 gr.
Ext. s. fldose { expectorant, 1-5 emetic \frac{1}{4}-1 dr.
Tr. sdose { expectorant, 5-30 emetic 1-4 dr.
Actions. Emetic, narcotic, expectorant. Use. Chronic bronchitis.
SANTONICA (Levant wormseed) 15-60 gr. Santonin
Actions. Anthelmintic for roundworm. Colors urine yellow, if acid; red, if alkaline. Produces xanthopsia. May produce toxic symptoms. **Uses. Roundworms.
SAPO (castile soap):
Empl. s. The for poisoning by a
Sapo mollis (soft or green soap).
Lin. s. raollis.
SARSAPARILLA.
Decott. S. Co
Ext. s. fl
- lantonin relieves the pains of crises of
is said ataxia of grown between
- 1/2 gr 2: Landburgeria

Ext. s. fl. co 1-1 dr
Syr. s. co 1-1 oz
Actions. Unknown.
Uses. Vehicle, chiefly.
SASSAFRAS 1-1 dr
S. medulla.
Mucilago s. m freely,
Oleum sassafras 1-5 M.
Actions. Like volatile oils.
Uses. Vehicle and flavoring agent.
and and and
SCAMMONIUM Cara 5-15 gr.
Res. s
Actions. Hydragogue cathartic-4 hrs. Anthel-
mintic for tape- and roundworm.
Uses. Obstinate constipation. Tape- and round-
Basis of Mediterraneon
SCILLA (squill) 1-5 gr.
Acetum s 10-45 M.
Ext. s. fl
Тт. s 5-30 Щ.
Syr. s 1-1 dr.
Syr. s. co. (hive syrup) 5-30 M.
Actions. Like digitalis. More of a gastro-intestinal
irritant. More powerful diuretic. Powerful ex-
pectorant.
Uses. Diuretic in chronic nephritis with digitalis.
In combination in chronic bronchitis.

THE ACTIONS AND USES OF DRUGS SCOPARIUS (broom) t dr. Ext. c. fl Action and use. Diuretic. SENEGA ... W.S...... 10-20 gr. Ext. s. fl...... 10-20 M. Syr. s..... 1-2 dr. Syr. scillee co. (hive syrup)..... 5-30 M. Actions. Ext. Irritant. Int. Gastro-intestinal irritant. On heart like digitalis. Expectorant. Diuretic. Use. Chronic bronchitis. *SENNA ... alvi I-3 dr. Confectio s..... 1-2 dr. Inf. s. co. (black draught) acts in 1-2 h, 1-2 oz. 5 Pulv. glycyrrhizæ co. (compound licorice - -powder) ... acts. in & h. 1-2 dr. (oil. Syr. s..... 1-1 oz. Actions. Purgative, acting on colon. Gripes. Yellow, watery stools. Urine red. Use. Purgative. Generally in combination. SERPENTARIA (Va. snakeroot)..... 10-30 gr. Ext. s. fl...... 10-30 ML Tr. s..... 1-2 dr. Action and use. Stomachic bitter. Prisoning dermet anzidiphil

SEVUM Preparatum

Uses. Making cerates.

SINAPIS ALBA (white mustard). NIGRA (black mustard).

Charta s.

OLEUM S. VOLATILE.

Linimentum s. co.

Actions. Ext. Irritant. Rubefacient. Counterirritant. Vesicant. Anodyne. Reflex cardiac and respiratory stimulation. Int. Stomachic. Emetic.

Uses, Ext. General counter-irritant and anodyne. Antemetic over stomach. Cardiac and respiratory stimulant. Int. Condiment. Emetic.

SODIUM:

Liquor sodæ..... 5-20 M. freely diluted.

Actions. Like liquor potassæ. Little used. Sodie carbonas mono hydratus

S. carbonas (sal sada, washing sada).... 5-15 gr. 8 " exsicentus 5-15 gr.

S. bicarbonas (baking soda, soda).... 5-60 gr.

Trochisci sodii bicarbonas, aa 3 gr. 1-6

Uses. Gastric sedative. Acid eructations. Dyspepsia.

S. phosphas..... 5 gr, to I oz,

S. sulphas (Glauber's salt)..... 1-8 dr.

S. phosphas exsic. 29r. 59. 59. 59. 59. 59. 59.

Potassii et sodii t rtras (purgative, 1-1 oz. (Rochelle salt)dose (diuretic 30-60 gr. Pulvis effervescens compositus (Scidlitz powder).
Actions. Saline purgatives. Sulphate most powerful. Sulphate and phosphate, cholagogues. Uses. Habitual constipation. Cholagogues. Dropsy. Ascites.
S. chloridum (common salt) 5-60 gr.
Uses. Food. Emetic (large doses). As rectal injections for oxyuris vermicularis.
S. sulphis
Uses. Sulphite used for gastric fermentation.
S. nitras ½-1 oz. Purgative.
S. acetas 10-60 gr.
Diuretic (little used). Used to make acetic acid, acetic ether, etc. S. chloras
Similar to potassium salt.
SODII ETHYLAS.
Uses. Mild caustic to remove warts, nævi, etc.
Rochelle Ralto 120 gr
nanels 40

	tion more prolonged	
86	SYLLABUS OF MATERIA MEDICA.	
S	ODII NITRIS 2-5 Spr. ætheris nitrosi (sweet spirit of nitre). 1-2	
	ctions. Int. Stomachic, carminative. Car stimulant. Vaso-dilator. Diaphoretic. Diur ses. Dyspepsia. Cardiac diseases with cede Febrile diseases. Nephritis.	
	ction and use. Antiseptic in gastric and intesfermentation.	
S	OMNAL 15-30	0 1
A	ction and use. Hypnotic.	
S	PARTEINÆ SULPHAS (fr. scoparius), 10-2	2 5
A	ction and use. Cardiac stimulant.	
S	PIGELIA (pinkroot)	
A	ction and use. Anthelmintic for roundworm.	
	PIRITUS GLONOINI (spirit of nitroglycerine)	

STAPHISAGRIA (stavesacre).
Action and use. Parasiticide for pediculi.

4 bitut asia naturalized in most countries THE ACTIONS AND USES OF DRUGS. STRAMONIUM (Jamestown werd, thornapple). S. semen.... 1-3 gr. Ext. s. series Y X X X 1 1 gr. HundExt. " " A 1-3 M Tr. " " 5-30 TL Ung. s..... 10% S. folia Stramonium. 1-5 gr. Actions. Like belladonna. Relaxes bronchial tubes more, It may make the heart irreg Uses. Chiefly asthma. Preparations made from the l STRONTIUM. S. lactas 1-2 dr. Actions. Diuretic. Poluces albumin in albuminuria. Sedative to heart Uses. Diuretje. Albuminuria from renal atrophy. Chronie rheumatism, gout, scrofula. Diabetes. Circhosis. Trop. africa *STROPHANTHUS. Tr. s. . (New Tr.) 2-10 Actions. Like digitalis. Feebler diuretic. 200 %. Uses. As for digitalis. Not so generally useful. STYRAX (storax, liquid amber)..... 5-20 gr. Actions. Like bals, of Peru and tolu. Use. Pediculi and scabies. is. from digitalis no dias

of. SULPHONAL & Sulphamenthenum 15-40 gr.

Use. Hypnotic-2 hrs. to act. Not depressing.

* SULPHUR:

S,	sublimatum (flowers of s.)	15-60 g.
**	præcipitatum (milk of s)	15-60 gr.
	lotum (washed s.)	
U	ng, sulphuris	30%

Actions. Ext. Parasiticide. Int. Mild laxative.
Uses. Ext. Scabies. Sulphur baths for rheumatism.
Acne. Int. Laxative for children and in piles and anal fissures. Slight constipation. Sometimes for skin diseases.

DEDESK MINISTER.

Calx	46	See	p. 27.	10-3
Sulphuris	iodidum.			I-4 gr.

Actions. Ext. Parasiticides. Int. Unknown.

Uses. Ext. Scabies. Chronic skin diseases. Baths for rheumatism, myalgia, etc. Int. Chronic rheumatism and skin diseases. Efficacy doubtful.

TABACUM (Lobacco).

497.

Nicotine.

Actions. Gastro-intestant irritation. Collapse.
Cardiac innervation disturbed. Motor nerves depressed. Respiration paralyzed. Myotic.
Uses. Rarely used. Poultice for epididymitis.

Hesos & Hesoy . . a severe initiant.

THE ACTIONS AND USES OF DRUGS. 89
TAMARINDUS freely.
Use. Mild laxative.
TXNACEXUM (tansy)
Action and use. Dangerous. Abortifacient.
TARAXACUM (dandelion)
TEREBINTHINA CANADENSIS (Canada bal-sam);
Action. Like turpentine. Use. As an adhesive varnish.
TERPINI HYDRAS 2-30 gr.
Actions. Antiseptic expectorant. Diuretic. Stimulates gu. mucous membrane. Uses. Acute and chronic bronchitis, pulmonary tuberculosis, whooping-cough. Chronic nephritis, cystitis, gonorrhea.

THEOBROMINÆ SODIO-SALIC	YLAS (diuretin), =
di methyl xanthine	15-30 gr.
Action and use. Diuretic.	
THERIACA (treacle, molasses):	
Actions. Demulcent. Slightly lax Uses. Flavoring agent, etc.	ative.
THYMOL 1200 Hac) 1-15 gr.
Actions and uses. Antiseptic and	parasiticide.
TRAGACANTHA: Mucilago t	indeterminate.
Uses. Demulcent, vehicle.	
TRIONAL = Sulphonethyl	nettamen 7-30 gr.
Action and use. Hypnotic. De	lirium tremens.
TRITICUM (couch grass)	1–8 dr. 1–8 dr.
Action and use. Diuretic.	
ULMUS (slippery elm):	7
Mucilago u	freely.
Use. Demulcent,	

	THE ACTIONS AND USES OF DRUGS. 91
	URETHANUM Sthyl Cardramate 5-30 gr.
1	Action and use. Hypnotic.
1	UVA URSI (bearberry)
	Actions. Diuretic. Astringent and disinfectant to urinary tract. Uses. Pyelitis, cystitis, gonorrhœa, etc.
1 9	VALERIANA 10-30 gr.
	Ext. v. fl 10-30 M.
	Tr. v ½-2 dr.
	Tr. v. ammon 1-2 dr.
	Ammonii valerianas 2-8 gr.
130	№ ; " 🗡 gr.
	Zinci Valiras +3 gr.
*	Actions. Of volatile oils. Reflex cardiac stimulant. Uses. Carminative. Fainting. Palpitation. Neuralgia. Hysteria. To hysteria was ammonisted by
	VANILLA: malingaless)
	Varieties Wenter Veneguela VENATRINA CONTRACTOR OF Alkaloide
	VERATRINA (mixture of alkaloids) 1 1 gr.
	Oleatum v
	Ung. v 4%
-	Valerian is supposed to be a 11 redative
-	only of value because of its assative anti-
nd	its bad taste

Actions. Ext. Irritant. Int. Gastro-intestinal irritant. Heart action greatly slowed. Motor and sensory nerves first stimulated, then paralyzed. Contraction of vol. muscle greatly prolonged.
Use, Ext. For neuralgia,
VERATRUM VARIOS (Am. hellebore) 1-5 gr.
Ext. v. v. fl 1-5 m.
Tr. VX 3 40%. 2-10 M.
Actions. Cardiac depressant.
Uses, As cardiac depressant, Puerperal eclampsia,
VIBURNUM OPULUS (crampbark) 1-2 dr. Ext. v. o. fl
Actions. Antispasmodic, tonic, diuretic. Uses. Nervous disorders of pregnancy. To prevent miscarriage. Spasmodic dysmenorrhæa. After- pains. Menorrhagia.
VIBURNUM PRUNIFOLIUM (black haw), 1-1 dr. Ext. v. p. fl
Actions and uses. See above.

WITELLUS (yolk of egg):

Uses. Emollient. Nutritive.

VOLATILE OILS.

Actions. Ext. Rubefacient, irritant, counter-irri-

tant, vesicant. *Int.* Stomachic, carminative; gastro-intestinal irritant in large doses. Expectorant. Diuretic. Stimulate g.-u. mucous membrane.

ZINCUM: .

*

Z. chloridum.

Liquor zinci chloridi.

1-10 m

Actions and uses. Ext. Caustic, antiseptic.

Z. sulphasdose { tonic emetic,	10-30	gr. gr.
Z. carbonas præcipitatus.		-
" oxidum Unguentum zinci oxidi. 2 % Oleatum zinci.	1-5	gr.
Z. acetastonic,		gr.

Actions. Ext. Astringent, hemostatic. Int. Direct emetic. Tonic.

Uses. Ext. General astringent. Int. Emetic. Diarrhœa. Hysteria, epilepsy, whooping-cough. Tonic.

ZINGIBER (ginger)	. 5-15 gr.
Ext. z. fl	. 5-15 M.
Oleores. z	$\frac{1}{2}-2$ Π .
Tr. z	.] -1 dr.
Syr. z	. 1 -2 dr.
TRANSPA	Walter.

Actions and uses. Like volatile oils.

ung, zinci stearatis 50 %. Contains

PREPARATIONS.

QUÆ:			
Aqua	anisi] -1	oz
"	aurantii florum fortior	indefin	ite
44	66 66	"	
"	ammoniæ fortior	3-6	π
• •	44	10-20	π
	amygdalæ amaræ	2-4	dr
"	camphoræ	1-2	οz
	chlori	1-4	dr
• •	chloroformi	1-4	đ٢
"	cinnamomi] -1	oz.
	creosoti	1-4	dr
• •	fœniculi	1 -1	oz
	hydrogenii dioxidi	1-3	dr
	menthæ piperitæ	1-2	oz.
	" viridis	-	oz
**	rosæ fortior	-	
**	44	**	

EXTRACTA:

Max	imum d	ose one grain or less:	
1-	Extract	um aconiti	10 18
2	**	belladonnæ foliorum	1 1 gr
3		cannabis indicæ	
4	44.	-conii	
5	166	digitalis	
6	-	nucis vomicæ	
7	a	opii	. 1-1 gr
8	44	physostigmatis	10 1 g
9	**	stramonii seminis	1-1 g
		ose more than one grain:	
E		n aloes	
		arniece radicis	
	**	cimicifugæ	
	**	einchonæ	5-30 gt
	**	colchici radicis	. 1-2 gr
		colocynthidis	. 1-2 g
	440	" compositum	. 5-15 gr
	***	ergotæ	- 5-15 gr
	**	euonymi	. 1-5 gr
		gentianæ	. 2-10 gr
		glycyrrhizæ	freely.
	11	hæmatoxyli	. 5-15 gt
	1.66	hyoscyami	. 1-2 gi
		-iridis	, 1-3 gr
	94	sialanos	

Ex	tractum	juglandia	5-30	gr.
	"	krameriæ	5–10	gr.
	"	leptandræ		_
	" .	- podophy tti	-	_
	44	quassiæ		
	4.6	rhei		
	" "	taraxaci		
	٠٠ ,	- uvo uroi		
			•	•
EXTR.	ACTA	FLUIDA:	•	
Mari	mum de	ose less than 10 minims:		
I		tum aconiti fluidum	l -2	π.
2	LAHACI	belladonnæ radicis fluid-	9 -2	ι.
_		um		π.
•	14	cannabis indicæ fluidum,	1-3	η.
3	"		2-5 1-8	•
4		capsici fluidum	2-8 ·	Π.
5 6		colchici radicis fluidum,		M.
_		colchici seminis "	1-5	π.
7		conii fluidum	2-5	π.
8		digitalis fluidum	1 -3	Π.
9		tobeliæ "	2–10	η.
10	**	nucis vomicæ fluidum	I - 4	η.
11	**	ipecacuanhæ fluidum,		
		dose { expectorant, emetic I	1-5 5-30	Π Π.
12		sanguinariæ fluidum,		
		dose { expectorant, emetic	I-5 1 -I	η. dr.

13	Extractum	scillæ fluidum	1-5	η.
14	44	stramonii seminis fluid-		
		um	1-3	η.
15	- 41	veratri viridis fluidum	1-5	η.
Maxi	mum dose l	between 10 and 30 minims	:	
1	Extractum	apocyni fluidum	2-20	η.
2	**	arniese radicis fluidum,	5-20	mt.
3	44	aromaticum "	10-30	η.
4	**	aspidospermatis "	5-30	η.
5	44	buchu fluidum	15-30	η.
6		calumbæ "	5-30	π.
7	44:	convallariæ fluidum		
8	10	cypripedii "	15-30	π.
9	16	eryodictyi "	15-30	щ.
IO	44	gelsemii fluidum		
II	0.	gentianæ "		
12	- 11	hyoscyami "	5-15	η.
13		iridfs "	10-30	m.
14	44	krameriæ "	5-30	щ.
15	40	lupulini "	100 100	
16		menispermi fluidum	5-30	π.
17	44	mezerei "		
18	169	phytolaccæ radicis fluid-	-	-
		um	5-30	Щ.
Iq	- 44	podophylli fluidum		-
20	**	sabinæ "		
21	**	senegæ "	1000	

PREPARATIONS.

22	Extract	ium serpeniariæ	; nuiuum.	• • • •	10-30	ա.
23	"	valerianæ	٠٠ .		10-30	Π.
24	"	xanthoxyli	٠٠ .		10-30	η.
25	"	zingiberis	" .		5-15	π.
Maxi	imum do	ose I drachm:				
Ex	tractum	calami fluidun	n] —I	dr.
	**	chiratæ ''			-	
		cimicifugæ flu			-	
	**	cinchonæ			_	
	"	cubebæ				•
	44	ergotæ] -1	
	4.4	eupatorii			-	
	4.4	geranii			1 -1	
	**	gossypii radic			•	dr.
		0 7.	15 11414411		-	-
		grindeliæ	**	• • • •] -1	
	**	guaranæ	"	• • • •] -1	-
		hydrastis		• • • •	•	-
	**	lappæ	4.6	• • • •	1 −1	
	4.6	leptandræ	**		} −1	dr.
	• •	pareiræ	"] -1	dr.
	**	pilocarpi	**		5–60	π.
		pruni Virginia	næ ''] —I	dr.
	**	quassiæ	"		} −r	dr.
	44	rhamni purshi	anæ fluid	um] —1	dr.
	**	rhei	**] —I	dr.
	**	rhois glabræ	**] _1	dr.
	**	rosæ	**] _1	dr.

Extractum	rumicis fluida	m		- -	dr.
44	sarsaparillæ f	luidum.] _I	dr.
**	** 1	Auidum	compo-		
	situm] -1	dr.
• •	scoparii-Auid	um		} - -	dr.
44	stillingiæ	• • • • •] -1	dr.
•	uvæ ursi	••		} −ı	dr.
"	viburni prun	ifolii flu	idum] –1	dr.
Maximum do	se more than	one drac	hm:		
Extractum	asclepiadis fl	uidum		1-2	d r.
4.4	aurantii	" .		1-2	dr.
**	castancæ	"		1-2	d r.
**	chimaphilæ	"		1-2	dr.
**	cocæ	٠٠		1-4	dr.
**	e usso-	٠٠		7-4	dr.
* *	dalcamaræ	٠٠		1-0	dr.
4.6	eucalypti	"		1-2	dr.
• 6	frangulæ	"	• • • • • • •	1-2	dr.
"	glycyrrhizæ	"		free	ly.
**	hamamelidis	"		1-2	dr.
"	matico	"		$\frac{1}{2}$ -2	dr.
"	rubi	"	· · · · · · · · ·	$\frac{1}{2}$ -2	dr.
**	scutellariæ	"		1-2	dr.
**	sennæ	"		1-3	dr.
**	spigeli.e	"		1-2	dr.
**	taraxaci	"	• • • • • • •	1-4	dr.
"	tritici	"		1-8	dr.
**	viburni opuli	٠٠	. 	1 -2	94.

LIQUORES:

CONTE			
Liquor	acidi arsenosi, 1% 5	-10	Π.
1.5	ammonii acetatis (spirit of Min-		
	dererus)	-8	dr.
**	arseni et hydrargyri iodidi (Dono-		
	van's solution), 1% each 1	-10	π.
	calcis (time-water) 1		
24	ferri acetatis 2		
4.6	" chloridi 2		
86	- eitratis 5		
**	" et ammonii acetatis (Bash-	-	100
	am's mixture) 1	-4	dr.
41 .	ferri nitratis 2	-	
**	" subsulphatis (Monsel's solu-	-	-
	tion) 2	-10	m.
44	ferri tersulphatis.	-	-
14	hydrargyri nitratis.		
11	iodi compositus (Lugol's solution),		
	I 5%, KI 10%	-10	т.
**	magnesii citratis		
11	plumbi subacetatis (Goulard's ex-		
	tract) e	xter	nal.
**	plumbi subacetatis dilutus (lead-		
	water) e	xten	nal.
11	potassæ, 5%		
	potassii arsenitis (Fowler's solu-	-	4.
	tion), 1%	2-10	m
	potassii citratis		1000
	bearings successive and a second second		-

Liqu	or sodæ, 5% 5-20	π.
"	sodæ chloratæ (Labarraque's solu-	
	tion) $\frac{1}{2}$ -I	dr.
11	sodii arsenatis (Pearson's solution),	
	1g 1-10	π.
"	so dii silicatis.	
**	zinci chloridi 1–10	η.
MASSA	E:	
Mass	sa copaibre	dr.
44	ferri carbonatis (Vallet's mass) 1-5	gr.
**	hydrargyri (blue mass, blue pill),	•
	33% 12-15	gr.
OLEA:		
Maxim	um dose 5 minims or less:	
	Oleum amygdalæ amaræ 1-1	П.
2	" anisi I-5	
3	" aurantii florum	
4	" betulæ volatile	
5	" cajuputi	π.
6	" cari 1-5	π.
7	" caryophylli	π.
8	" cinnamomi	π.
9	" coriandri 2-5	
10	" fœniculi	Π.
11	" gaultheriæ	η.

12	Oleum	hedeomæ 1-5 M.
13	44	lavandulæ florum 1-5 M.
14	**	menthæ piperitæ 1-5 M.
15	**	" viridis 1-5 M.
16	16	myristicæ 1−3 ៕.
17	**	phosphoratum, 17 1-5 M.
18		pimentæ 1-5 M.
19	45	rosmarini 1-5 M.
20	15	sabinæ 1-5 M.
21	184.1	sassafras 1-5 M.
22	**	sinapis volatile 1-1 m.
23	**	thymi 1-5 M.
24	11	tiglii (croton vil) 1-2 m.
Maxi	mum d	ose between 5 and 30 minims:
1	Oleum	chenopodii 2-10 Π.
2	**	copaibæ 5-15 m.
3	44 4	cubebæ 5-20 M.
4	25 1	erigerontis 5-15 M.
5	990	eucalypti 5-30 M.
6	44	juniperi 2-10 M.
7	44	santali 2-10 M.
8	44	terebinthinæ 5-30 M.
9	- 11	" purificatum 5-30 П.
Max	imum d	ose more than one drachm:
Ol	eum an	nygdalæ expressum 1-4 dr.
	11 10-1	i (linseed) 1-2 oz.
	lin	11 (unseed)
	IIII	orrhuæ (cod liver)

Oleum	olivæ	freely.
4.6	ricini (castor)	1 -2 oz.
**	sesami	-
Oleum	aurantii corticis.)
4.4	bergamottæ.	
	limonis.	} flavoring.
**	rosæ.	}
Oleum	adipis.	used in pharmacy,
"	æthereum.	sed in pharmacy,
Oleum	mayaroite	perfume,
**	gossypii seminis.) -
"	picis liquidæ.	
"	cadini.	externally,
44	theobromatis.	
OLEORES	CINIÆ.	,
Oleore	sina aspidii	-
.,	capsici	-
		····· 5-30 叽.
"	lupulini	- •
"	piperis	
"	zingiberis	····· ½−2 ៕.
PILULÆ	:	
Pilulæ	aloes, 2 gr	1-4 pills.
**	" M searchaide. Di	
"	" ferri, I gr	

Elever of gr. L. layativas Co. J. Struck. 125 " > 2 kill lever selens 1."
104 PREPARATIONS.
Pilulæ aloes et mastiches ("Lady
Webster"), 2 gr 1-3 pills.
" aloes et myrrhæ, 2 gr 2-10 "
" antimonii compositæ (Plum-
mer's pills)
" asafœtidæ, 3 gr 1-4 "
" catharticæ compositæ 1-3
Calomel
Pilulæ catharticæ vegetabiles 1-5 Ext. colocynthidis co
Pilulæ ferri carbonatis (Blaud's pills). 1-2 "
" " iodidi
" opii, I gr
" phosphori, 100 gr 1-5
" rhei, 3-gr
" compositæ, 2 gr 1-3 "
PULVES:

r uivis	-antimonians (James powder)	3-15 gr.
64	aromaticus	10-30 gr.

Pulvis c	retæ compositus	5-00	gr.
" e	ffervescens compositus (Seidlitz		
	powder) one	powd	er.
" g	lycyrrhizæ compositus		
" ij	pecacuanhæ et opii (Dover's		
	powder), each 10%	3-15	gr.
" j	alapæ compositus] - 1	dr.
" n	norphinæ '' (Tully's pow-		
•	der), 8 gr. to 1 oz	5-15	gr.
" r	hei compositus (Gregory's pow-		
	der)] —I	dr.
RESINÆ:			
Resina.			
**	eepaibæ	5-15-	gr.
	jalapæ		
	podophylli] -1	gr.
"	scammonii	1– 8	gr.
SPIRITUS	: _		
Maximum	dose one drachm or less:		
Spiritus	ætheris] —I	dr.
"	" compositus (Hoffman's		
	anodyne)	5-60	η.
"	ammoniæ I	0-60	η.
"	amygdalæ amaræ 1	5-45	η.
"	camphoræ	} −1	dr.

Spi	ritus c	chloroformi (chloric ether)	1-1 dr.
	(cinnamomi	10-30 €.
		glonoini (spirit oj nitroglyce-	
		rine)	1-3 П.
14		juniperi	1-1 dr.
4		avandulæ	1-I dr.
9	. 1	limonis	flavoring.
		menthæ piperitæ	5-15 M.
	11	" viridis	5-15 M.
	,	myristice	1 dr.
		phosphori	8-40 III.
Monie		dose more than one drachm:	- 4
		Court burde a strong auto account.	- 6
1	Spirit	us ætheris nitrosi (sweet spirit	
		nitre)	Contract of the Contract of th
2		ammoniæ aromaticus	
3	**	anisi	
4	**	aurantif	vehicle.
5	45	compositus	San San San San
6	**	frumenti (whiskey)	indefinite.
7	**	gaultheriæ	1-2 dr.
8	44	juniperi compositus	
9	**	myreire	perfume.
10	- 11	vini gallici (brandy)	indefinite.
SYRU			
Syn			
	**	acaciæ	"

Syrupus	acidi citrici	1-4 dr.
11	" hydriodici	1-1 dr.
110	allii	r-4 dr.
14	althouse	freely.
44	amygdalæ	44
16	aurantii	vehicle.
46	" florum	freely.
- 4.1	calcii phosphatis	1-2 dr.
0	The state of the s	15-60 M.
4.6	ferri iodidi	5-30 TL.
4.6	" quininæ et strychninæ	
	phosphatum	1-1 dr.
**	hypophosphitum	1-2 dr.
188	" cum ferro	1-2 dr.
**	ipecacuanhæ 5	-2 dr.
44	krameriæ	1-4 dr.
**	lactucarii	1-4 dr.
1.6	picis liquidæ	1-4 dr.
34	pruni Virginianæ	1-4 dr.
44	rhei.	-
84	" aromaticus ı dr.	for child.
44	rosæ	1-2 dr.
46	rubi	1-2 dr.
- 41	u idei,	freely.
-	sarsaparillæ compositus	1-I oz.
- 11	scillæ	1-1 dr.
44	" compositus (hive syrup)	5-30 TL

PREPARATIONS.

Syr	upu	s senegæ	1-2	dr.
		sennæ] —1	oz.
	"	tolutanus] —I	oz.
	••	zingiberis	1-2	oz.
TINCT	UR.	Æ:		
Maxir	nun	n dose half a drachm or less:		
1	Tr.	aconiti	1-5	π.
2	"	aconiti	5 TO	-π.
3	• •	belladonnæ foliorum	5 -3 0	π.
4	"	cannabis indicæ	5-30	π.
5	"	cantharidis	1-15	π.
6	"	colchici seminis,	10-30	π.
7		digitalis	5-30	
8	"	iodi	1-5	η.
9	"	ipecacuanhæ et opii	3-15	Π.
10	"	lobeliæ	5-30	π.
11	"	nucis vomicæ	5-20	π.
12		opii (laudanum)	3-20	π.
13	"	" camphorata (paregoric), for		
		child	3-20	η.
. 14	**	opii deodorata	3-20	π
15	"	physostigmatis	10-30	π
16	"	sanguinariædose expectorant, emetic	5-30 1-4	η. dr.
17	**	scillæ	5-30	
18	"	stramonii seminie	5-30	

PREPARATIONS.		109
19 Tr. strophanthi	2-10	π.
20 " veratri viridis		
Maximum dose one drachm:		
Tr. aloes	15-60	π.
" asafœtidæ	30-60	-
" arnicæ florum.	15-45	-
" benzoini	30-60	
" composita	30-60	
" capsici	5-60	-
" ferri chloridi	5-60	-
" gelsemii	15-60	-
" guaiaci	30-60	-
" " ammoniata	30-60	-
" lavandulæ composita	30-60	-
" moschi	15-60	-
" myrrhæ	_	-
" zingiberis	•	-
zingiberis	15–60	··L.
Maximum dose more than one drachm:		
Tr. aloes et myrrhæ	. I-21/2	dr.
" antiperiodica (Warburg's tincture)	. I-4	dr.
" aurantii amari	. I-2	dr.
" " dulcis	. I -2	dr.
😭 bryoniæ	. I-5	dr⅓
" calendulæ		dr.
" calumbæ	•	dr.
" cardamomi	•	dr

Γr.	cardamomi composita		
"	catechu "	1-2	dr.
"	chiratæ	1-2	dr.
"	cimicifugæ		
"	cinchonæ	1 -2	dr.
••	" composita	I-4	dr.
"	cinnamomi	I-2	dr.
"	croci	1-2	dr.
"	cubebæ	1-5-	dr.
"	gallæ		
٠.	gentianæ composita		
"	humuli	•	
41	hydrastis		
	hyoscyami	_	
٠,	kino	•	
• •	krameriæ	-	
	lactucarii	-	-
"	matico	-	
	opii camphorata		
	pyrethri.		
"	quassiæ	l -2	dr.
	quillajæ		
"	rhei		
	" aromatica	•	
	44 dulcie.	_	
	serpentariæ	-	-
	sumbul		

furine SOy ratine alus unbarle

PREPARATIONS.	trr
Tr. tolutana	½-2 dr. ½-2 dr.
VINA:	
Vinum album	. menstruum.
" antimonii	5-60 III.
" colchici radicis	-
" " seminis	
" ergotæ	• .
" ferri amarum	•
" " citratis	•
" ipecacuanhæ	
" opii (Sydenham's laudanu	
10%	3-20 ¶.
" rubrum	. menstruum.

A CLASSIFICATION OF DRUGS, BASED CHIEFLY ON PHYSIO-LOGICAL ACTION.

	 A drug or remedial agent which pro- y affects nutrition.
Mercu	ry.
Iodine	and its compounds.
Arseni	c
Colchi	cumradix 2-8 gr., semen 1-5 gr.
Phospl	orus and the hypophosphites.
Organi	c extracts, e.g., thyroid, etc.
porary	tic. A substance capable of producing tem- loss or impairment of feeling or sensation, are divided into general and local.
	General.
Chloro	form 2−20 ¶.
Ether.	5-60 π.
Nitrou	s oxide.
Ethyle	ne bichloride.
Methy	ene ''
Pental.	

Local.

Cold (ice or rapidly evaporating liquids). Cocaine
Analgesic. A medicine, drug, or remedial agent which brings about an incapacity of feeling pain in a part, although the tactile sense may be more or less preserved.
Applied especially to:
Antipyrin 3-20 gr.
Phenacetin 5-10 gr.
Antifebrin 2-5 gr.
Exalgin ½-3 gr.
And other coal-tar products which are not official.
Anaphrodisiac. A drug or remedial agent capable of depressing sexual desire.
Bromides.
Opium 1-2 gr.
Potassium iodide 5-60+ gr.
Belladonnafolia 1-5 gr.
Hyoscyamus 5-15 gr.
Stramoniumsemen 1-3 gr., folia 1-5 gr.
Anhidrotic. A drug which checks or tends to check sweating.
Atropine (belladonna), 120 ng gr.
Hyoscyamus 5-15 gr.

Stramoniumsemen 1-3 gr., folia 1-5 gr.	
Agaricin 1 2 gr	
Acids.	
Picrotoxine	
Muscarine 1/8-2 gr	
Nux vomica 1-4 gr	
nodyne. A drug or remedial agent capable of re	Į
lieving pain.	
Cold. Local Anodynes.	
Aconite ½-2 gr.	
Menthol	
Carbolic acid 1-1 gr.	
Iodoform	
Chloral 5-20 gr.	
Dilute hydrocyanic acid 1-3 19.	
Sodium bicarbonate 5-60 gr.	
Belladonnafolia 1-5 gr.	9
Stramoniumsemen 1-3 gr., folia 1-5 gr.	
Hyoscyamus 5-15 gr.	
Opium 1-2 gr.	
Veratrinedo 10 gr	
Zinc oxide 1-5 gr	
Certain volatile oils.	

Antacid. An alkali used as a remedy for acidity in the stomach.

Ammonia.

Potassium (some preparations).	
Sodium " "	
Calcium "	
Magnesia 5-6	gr.
Magnesia ponderosa 5-6	gr.
Anthelmintic. A drug or remedial agent use	d for
the removal of intestinal worms. Taniafa useful against the several varieties of tapeworn	
For Tapeworm.	
Aspidium 1-1	dr.
Granatum ‡-1	dr.
Kousso 2-4	dr.
-Kamala	dr.
Pepo 1-3	OZ.
Turpentine 1-4	dr.
Chloroform) III.
For Roundworm.	
Santonica	gr.
Spigelia	dr.
Senna 1-3	dr.
Chenopodium 15-3	gr.
For Threadworm.	
Mostly as injections:	
Tannic acid 1-2	ogr.
and drugs containing it.	

Turpentine 5-30 Π., ½-4 dr.
Sodium chloride 5-60 gr.
Carbolic acid ½-1 gr.
Vinegar.
Ferric salts.
Soapsuds.
Vegetable bitters - columbaste
Antiperiodic. A drug for the cure of diseases exhibiting periodicity, especially malarial fever and its manifestations.
Quinine, 1-20 gr. Warburg's tincture, 1-4 dr. Cinchona 10-60 gr,
Arsenic
Eucalyptus 1-2 dr.
Hydrastis 5-60 gr.
Salicin 5-30 gr.
Salicylic acid 5-60 gr.
Oleum gaultheriæ
Methyl salicylate 1-5 η.
Berberine.
Antipyretic. A drug or remedial agent which diminishes the body temperature.
Cold.
Evaporating liquids.
Antipyrin 3-20 gr.
Antifebrin 2-5 gr.

Carbolic acid...... 1 gr.

An

CLASSIFICATION OF DRUGS.

Hydrogen peroxide	1-3	đr.
Potassium permanganate	1-2	gr.
Boric acid		-
Iodoform	1-3	gr.
Iodine.		•
Sulphurous acid	1-2	dr.
Lime.		
Eucalyptus	1-2	dr.
Creolin	1-5	π.
Lysol.		
Chlorine.		
Bromine.		
Benzoin.		
Zinc chloride.	•	
Zinc oxide	1-5	gr.
Ferric oxide.		_
Alcohol.		
Menthol] -2	gr.
Balsam of Peru		
Balsam of tolu	10-30	gr.
Quinine	1-20	gr.
Intestinal Antiseptics.		
Creosote	1-2	111 .
Carbolic acid		
Salol		
Salicin		-

CLASSIFICATION OF DRUGS.	119
Salicylic acid	5-60 gr.
Bismuth salicylate	5-20 gr.
Resorcin	2-5 gr.
Boric acid	5-15 gr.
Eucalyptus	1-2 dr.
Thymol.	1-15 0
Sodium hyposulphite	5-20 gr.
" sulphocarbolate	5-30 gr.
Sulphurous acid	1-2 dr.
Naphtol	1-20 gr.
Naphtalin	1-20 gr.
Corrosive sublimate	1 1 gr.
	10.0
tispasmodics. Agents which arrest	or cure
spasms or convulsions. Applied espessasm of bronchial tubes.	ecially to
Stramoniumsemen 1-3 gr., folia	1-5 or
Belladonnafolia	
Hyoscyamus	
Grindelia	
Aspidosperma	
Ether	0
Chloroform	-
Opium	2 22 22
Chloral	-
Amyl nitrite	- 0
***************************************	-3 m

Anti

Cannabis Indica. 2-5 gr. Conium. 2-5 gr.
Aphrodisiac. A drug which increases the sexual desire.
Strychnia
Cantharides, tr 1-15 M.
Cannabis Indica 2-5 gr.
Camphor
Phosphorus
Damiana $\frac{1}{2}$ -I oz.
Astringent. A drug which contracts the tissues and vessels of the body, thereby diminishing or checking excessive discharges.
Mineral.
Silver salts.
Lead ''
Zinc "
Copper salts.
Ferric "
Alum 5-30 gr.
Bismuth salts.
Sulphuric acid.
Calcium (some preparations).
Lime-water 1-8 dr,

Vegetable.

(Galla, tr	I-2	dr.
	Quercus alba.		
	Krameria	5-30	gr.
	Kino	5-30	gr.
Tannic acid.			•
	Hæmatoxylon, ext	5-15	gr.
	Cinnamon.		
	Eucalyptus gum Hamamelis, ext. fl	2-10	gr.
			dr.
	• • • • • • • • • • • • • • • • • • • •		
	folia		gr.
• •	•••••	5-[15	gr.
Stramonium	semen 1-3 gr., folia	1-5	gr.
	ant or Sedative. A drorce or frequency, or both he heart.		
Aconite		. 1-2	gr.
	e. 		
Antimony.	•		
Dilute hydrocy	anic acid	. 1-3	η.
Cardiac Stimula force or freque the heart.	nt. A drug which incrency, or both force and freq	eases Iuency	the , of
Digitalis		1-3	gr.

Ammonia.	
Alcohol.	
Caffeine	1-5 gr.
Ether	5-60 m.
Nux vomica	1-4 gr.
Strophanthus, tr	2-10 Ⅲ.
	1-2 gr.
	1-1 gr.
	1-3 щ.
Convallaria, ext. fl	5-30 TL.
Sodium nitrite	2-5 gr.
Nitroglycerine	100 au gr.
	1-5 gr.
	10-30 Щ.
The state of the s	I-20 gr.
The state of the s	1-2 gr.
Arsenic	10 10 gr.
Mustard, externally.	00 10 8
	Alcohol. Caffeine. Ether. Nux vomica. Strophanthus, tr. Spartein, sulph. Adonidin. Amyl nitrite. Convallaria, ext. fl. Sodium nitrite. Nitroglycerine. Squill. Cereus grandiflorus, ext. fl. Quinine. Opium. Arsenic.

Carminative. A drug which causes the expulsion of gas from the stomach and intestines.

Stomachics, q. v.

Bitters.

Condiments.

Podon	Hyllum			1.00	5-20
					1-10
					2-8
	nth				1-2
	mus				5.5
					1-5
	n phospha			200	
Ipecac	uanha	******	1-2	gr.,	15-30
Dilute	nitric acid	1			5-30
Dilute	nitrohydre	ochloric a	cid		5-30
	ive sublim				1 1
	n benzoate				5-60
**					1-8
Potassi	ium "				1-4
	cum				1-5
Ammo	nium benz	coate			5-20
Rhuba	rb				10-30
					5-20
1	ony				5-15
	C				1 1

to the skin with the object of altering the size of the vessels of the subjacent viscera. (See Irritants.)

Deliriant. A drug which produces a state of mental agitation, characterized by restlessness, incoherence of speech, and various sensory illusions.

And best .. Bile (04 years)

Salinglates quimer aliabet

Indoform

124

CLASSIFICATION OF DRUGS.

Belladonnafolia	1-5	gr.
Stramonium semen 1-3 gr., folia	1-5	gr.
Hyoscyamus	5-15	gr.
Cannabis Indica	2-5	gr.

Demulcent. A drug or remedial agent which protects and soothes a part, especially a mucous membrane,

Flaxseed.

Glycerine.

Gelatine.

Gum.

White of egg.

Isinglass, etc.

Diaphoretic. A drug or remedial agent which increases the amount of perspiration.

Warmth, externally.

Hot drinks.

Pilocarpus	5-60 gr.
Alcohol.	-
Dover's powder	3-15 gr.
Sweet spirit of nitre	1-2 dr.
Ammonium acetate, liq	2-8 dr.
Potassium citrate	5-30 gr.

acetate.....

5-60 gr.

Antimony.

Nicotine.

Ipecacuanha 1-2 1	gr.,	15-30	gr.
Opium		1-2	gr.
Camphor		3-20	gr.
Senega		10-20	gr.
Cubeb		30-60	gr.
Colchicumradix 2-8 gr., ser	nen	1-5	gr.
Quinine		1-20	gr.
Salicin		5-30	gr.
Lobelia		1–10	gr.
			•
Diuretic. A drug or remedial agent w	hich	increa	ses
the amount of urine excreted.			
Water.			
Liquor potassæ		5-30	π.
Potassium acetate		5–60	gr.
" citrate		5-30	-
'' nitrate		5-20	gr.
" bicarbonate		5–60	gr.
Sodium acetate		10-60	-
Lithium.			•
Sweet spirit of nitre		1-2	dr.
Nitrites.			
Digitalis		1-3	gr.
Scoparius] —I	_
Alcohol.		-	
Strophanthus, tr		2-10	π.
Diuretin			•
Calomel (small doses).			•
\			

Caffeine	1-5 gr.
Squill	1-5 gr.
Cantharides, tr	1-15 M.
Buchu	15-30 gr.
Juniper (also in gin).	
Turpentine 5-30 M,	} −4 dr.
Uva ursi	l −1 dr.
Cubeb	30-60 gr.
Copaiba	1 −1 dr.

Ecbolic. See Oxytocics.

Emetic. A drug or remedial agent capable of producing vomiting.

Acting on the Stomach.

Warm water.	
Sodium chloride	5-60 gr.
Alum	5-30 gr.
Ammonium carbonate	2-15 gr.
Copper sulphate $\frac{1}{8}$ gr.,	2-20 gr.
Zinc " $\frac{1}{2}$ -2 gr.,	10-30 gr.
Tartar emetic $\frac{1}{10}$	
Turpeth mineral	s + gr .
Ipecacuanha $\frac{1}{2}$ -2 gr.,	15-30 gr.

Acting Centrally.

Apomorphine	1 ₅ -1	gr.,	$\frac{1}{20} - \frac{1}{10}$	gr.
Fartar emetic		10-	-1-2	gr.
pecacuanha	1-2	gr.,	15-30	gr.

Donogui	20 20 8
Squill	1-5 gr.
Zinc sulphate $\frac{1}{2}$ gr.,	
Copper " $\frac{1}{8}$ gr.,	2-20 gr.
Antemetic. A drug or remedial agent which or tends to relieve vomiting.	ch relieves
Acting Locally.	
Ice.	
Carbonated waters.	
Very hot water.	
Bismuth subnitrate	. 5–60 gr.
" subcarbonate	. 5–60 gr.
Opium	. 1 -2 gr.
Dilute hydrocyanic acid	. 1−3 ៕.
Cocaine (hydrochlor.)	.] -2 gr.
Cereum oxalate	. 1–8 gr.
Tincture of iodine	. 1−5 M.
Wine of ipecac	. 1-60 П.
Menthol	. 1 -2 gr.
Carbolic acid	. <u>1</u> -1 gr.
Creosote	. } −2 ∏.
Chloroform	. 2−20 M.
Ether	. 5-60 П.
Acting Centrally.	
Opium	1-2 gr.
Bromides.	- 0
Chloral	2-30 &s

Amyl nitrite 1-3 M. Nitroglycerine 100 no gr. Dilute hydrocyanic acid 1-3 M.
Dilute hydrocyanic acid 1-3 m,
A11-1
Alcohol,
Hyoscyamine (sulph.) 1 1 1 1 gr.
Emmenagogue, A drug which increases or restores the menstrual flow.
Apiol 5-15 gr.
Potassium permanganate 1-2 gr.
Manganese dioxide 2-10 gr.
Aloes ½-10 gr.
Cantharides, tr 1-15 M.
Myrrh 5-30 gr.
Rue, oleum 14 m.
Tanny t dr.
Sabine 5-15 gr.
Pennyroyal, oleum 1-5 π.
By Improving Nutrition.
Tonics.
Iron.
Arsenic 1 1 gr.
Copper,

Emollient. An agent which softens and protects parts, especially the skin.

Hot water.

Poultices Bland oils, etc.

Escharotic or Caustic. An agent which destroys the vitality of the part to which it is applied.

Heat.

Sulphuric acid.

Nitric

Chromic

Mercuric nitrate.

Caustic potash.

soda

Arsenic $\frac{1}{40-10}$ gr.

Zinc chloride.

Sodium ethylate.

Expectorant. A drug or remedial agent which aids or causes the expulsion of the bronchial secretion. They are stimulating or depressing, according to their action on the heart.

Stimulating.

Ammonium salts.

Senega..... 10-20 gr. Squill...... 1-5 gr. Benzoic acid..... 5-15 gr. Benzoin.

Balsam of tolu...... 10-30 gr. " Peru..... 10-30 ¶.

Sulphur..... 15-60 gr.

Depressing.

Antimony.
Ipecacuanha ½-2 gr., 15-30 g
Apomorphine
Lobelia
Pilocarpus 5-60 g
Alkalies.
Potassium iodide, 5-60+ gr
Galactagogue. A drug which increases the quantity of milk secreted.
Pilocarpus 5-60 gr
Castor-oil leaves
Alcohol.
Antigalactagogue. A drug which diminishes the quantity of milk secreted or arrests the secretion.
Belladonnafolia 1-5 gr
Hematinic. A drug which increases the amount of hemoglobin in the red blood cells.
Iron.
Arsenic 10 10 gr
Potassium permanganate.
Copper.
Hemostatic. An agent which stops bleeding.
Cold.
Most astringents q. v.

Calcium chloride, in bleeders 5-20 gr. Ergot 30-60 gr.
Hypnotic. A drug which produces sleep closely resembling natural sleep.
Opium. \$\frac{1}{4} - 2\$ gr. Chloral 5-20 gr. Butyl chloral 5-20 gr. Chloralamide. 10-30 gr. Bromides. 5-60 gr. Monobromated camphor. 2-10 gr. Hyoscyamus 5-15 gr. Cannabis Indica 2-5 gr. Sulphonal 15-40 gr. Trional 7-30 gr. Paraldehyde \$\frac{1}{4} \text{ dr.} Urethane. 5-30 gr. Somnal 15-30 fl. Amylene hydrate \$\frac{1}{4} \text{ dr.} Humulus, tr 1-2 dr. Lupulin 5-30 gr. Lactucarium 5-60 gr. Alcohol.
Irritant. An agent which causes dilatation of the vessels of the part to which it is applied—generally accompanied by pain. Heat. Aqua ammoniæ

Mercuric nitrate.
Arsenous acid
Tartar emetic $\frac{1}{10}$ - $\frac{1}{2}$ - 1 -2 gr.
Antimony chloride.
Silver nitrate 1-1 gr.
Zinc chloride.
Iodine.
Bromine.
Chlorine.
Phosphorus 100 gg.
Cantharides, tr I-15 M.
Croton oil $\frac{1}{4}$ π .
Copper nitrate.
" sulphate ½-½ gr., 2-20 gr.
Volatile oils.
Chrysarobin.
Camphor 3-20 gr.
Capsicum 1-8 gr.
Senega 10-20 gr.
Ipecac ½-2 gr., 15-30 gr.
Mineral acids.
Carbolic acid ‡-1 gr.
Creosote $\frac{1}{2}$ \mathfrak{m} .
Alcohol.
Chloroform 2-20 ¶. confined.
Ether 5-60 π.)
Burgundy pitch.
Canada "

Mydriatic. A drug which dilates the pupil.

Atropine (sulph.)	120 gr.
Homatropine (hydrobrom.)	120 20 gr.
Hyoscyamine (sulph.)	1 1 gr.
Daturine.	
Cocaine (hydrochlor.)	1-2 gr.
Gelsemine	1 0 20 gr.
Muscarine	1-2 gr.
Angethetics (late)	

Myotic. A drug which contracts the pupil.

Physostigmine (sulph.)	100 30 gr.
Opium	1-2 gr.
Pilocarpine (hydrochlor.)	1-1 gr.
Nicotine.	

Anæsthetics (early).

Narcotic. A drug which causes an unequal disturbance of the functions of the brain (Brunton). A substance which not only produces sleep, but also in large doses depresses the functions of respiration and circulation (White).

General anæsthetics.

Opium	1-2 gr.
Chloral	5-20 gr.
Belladonnafolia	1-5 gr.
Hyoscyamus	5-15 gr.
Stramonium folia 1-5 gr., semen	1-3 pr

Alcohol.	
Cannabis Indica	2-5
Lupulin	5-30
Oxytocic. A drug which increases uterine conduring or immediately after labor.	traci
Ergot 30)–6 0
Tansy	⊢ı
Rue, oleum	-5
Sabine	5 -/ 15
Cotton-root bark	⊢r
Quinine 1	-20
Hydrastis 5	-60
Purgatives are subdivided as follows:	
Laxatives, which promote the action of the by slightly stimulating peristalsis:	bow
Brown bread, etc., containing cellulose resi	due.
Sugars.	
Fruits, e.g.,	
Prune,	
Fig,	
Apples (with skin).	
Cassia 1	-8
Olive oil f	reely
Castor oil (in small doses).	
Glycerine 5	–60
Sulphur	-6 0

Magnesium.			
Nux vomica	1-4	gr.	
Belladonnafolia	1-5	gr.	
Hyoscyamus	5-15	gr.	
Stramoniumsemen 1-3 gr., folia	1-5	gr.	
Simple Purgatives, which promote the act bowels by stimulating peristalsis and i the secretion.			
Rhubarb 1-5 gr.,	10-30	gr,	8-
Aloes	1-10	gr.	24
Cascara sagrada	1-1	dr.	9
Senna	1-3	dr.	-
Ox-gall	5-15	gr.	
Drastic Purgatives or Cathartics, which a powerful than simple purgatives, and a tively stimulating peristalsis and the secretion.	ct by	ac-	
Calomel	1-10	gr.	
Colocynth	2-8		
Podophyllum	5-20	gr.	
Aloes	1-10	-	
Gamboge	1-5	~	
Turpentine 5-30 nl.,	1-4	720	
Croton oil	1-2	η.	
Hydragogue Catharties, or catharties which secretion of large quantities of liquid.	cause	the	
Elaterin	1 1 20 12	gr	

Jalap				
Scammony				5-15 g
Saline Purgatize tracting liquid liquid stimu	id from th	he blood,		
Magnesium su				
Sodium	**			. 1-8 d
" phosp	hate		5	grI o
Potassium and				
" sul	phate	******		1-4 d
Cholagogue Purg	gatives.	See Choi	LAGOGUES	
ing vigor; a f cruits the vital	ood, core	lial, or m	edicine w	
Examples:				
Cod-liver oil .				1-4 d
Iron.				
Quinine				1-20 g
Vegetable bitt				
Rubefacient. A				

Rubefacient. An agent which produces redness of the skin, at the point of application, through vascular dilatation.

See IRRITANT.

Sedative. A drug or remedial agent which pos-

sesses	the	property	of	allaying	irritation,	irrita-
bility,	or p	ain.				

Sialagogue. A drug or agent which increases the secretion of saliva.

Acting on Nerve-endings.

Pilocarpus		
Muscarin	}- 2	gr.
Iodine.		
Mercury.		
Tobacco.		
Physostigma	1-2	gr.
Acting Reflexly.		
Acids.		
Acid salts.		
Sapid substances.		
Alcohol.		
Chloroform	2-20	π.
Ether	5 – 60	π.
Emetics.		
Antimony.		
Ipecacuanha ½-2 gr., 1	5-30	gr.
isialagogue. A drug or agent which din	ninisl	hes

Antisialagogue. A drug or agent which diminishes the secretion of saliva.

Acting on Nerve-endings.

Belladonna.....folia 1-5 gr.

Ipecacuanha ½-2 gr.,	15-30 gr.
Lobelia	1-10 gr.
Tobacco.	_
• Hydrocyanic acid (dil.)	1-3 M.
Quinine	
Veratrine	

VARIOUS GROUPINGS.

DRUGS WHICH MAY PRODUCE ERUPTIONS.

Aloes] -10	gr.
DRUGS EXCRETED IN THE MI	LK.	
Turpentine 5-30 ¶,	1 –4	dr.
Sulphonal	15–40	_
Salicylic acid	5–60	•
Quinine	1-20	_
Phenacetine	5–10	_
·Opium	} -2	_
· Iodides.		
Cubeb	30–60	gr.
Copaiba] -1	
Chloral	5-20	•
- Bromides	5–60	•
.Belladonnafolia	1-5	_
, Antipyrin	3-20	_
• Acetanilide	2-5	gr.
- Arsenic	.go−7,º.	gr.

-Antimony.

· Bromides	5-60 gr.
Castor oil.	
Copaiba	
Garlic	15-30 gr.
· Iodides.	
· Iron.	
Jalap	5-20 gr.
· Lead.	
Mercury.	
*Opium	
·Quinine	I-20 gr.
	I-20 gr.
· Quinine	I-20 gr. IO-30 gr. 5-I5 gr.
Quinine Rhubarb I-5 gr., Scammony. Senna.	I-20 gr. I0-30 gr. 5-I5 gr. I-3 dr.
· Quinine	I-20 gr. I0-30 gr. 5-I5 gr. I-3 dr.
Quinine Rhubarb I-5 gr., Scammony. Senna.	I-20 gr. I0-30 gr. 5-I5 gr. I-3 dr. I5-60 gr.
Quinine Rhubarb I-5 gr., Scammony. Senna Sulphur.	I-20 gr. I0-30 gr. 5-I5 gr. I-3 dr. I5-60 gr.

DRUGS WHICH AFFECT THE COLOR OF THE FECES.

Iron-black.

Bismuth-dark slate color or black.

Calomel-spinach-green.

Logwood-red.

All cholagogues render the feces darker.

Gastro-intestinal irritants may cause the appearance of blood.

Semma - Red

DRUGS AFFECTING COLOR OF URINE.

- Carbolic acid—dark from presence of pyrocatechin and hydrochinon.
- -Diuretics all lower color.

Hæmatoxylin (logwood)-reddish or violet.

- Potassium chlorate paroxysmal hemoglobinuria (sometimes).
- Renal irritants—dark from concentration or blood (hematuria).

Rhubarb-yellow.

Santonin—greenish-yellow or saffron, if acid; purplish red, if alkaline.

Senna-reddish.

DRUGS WHICH MAY CAUSE THE URINE TO REDUCE FEHLING'S SOLUTION AND TO RESPOND TO SOME OTHER TESTS FOR SUGAR.

Acidum	arsenosum	1 1 gr.
44	hydrocyanicum dilutum	1-3 П.
44	salicylicum	5-60 gr.
11	sulphuricum dilutum	
Alcohol.		

5-20 gr.

2-20 ♏.

<u>∔</u>-ւ dr.

Chloral

Chloroformum.....

Copaiba

Glycerinum Mercury, salts of. Morphia, " " Strychninæ sulphas Oleum terebinthinæ 5-3	1 1 gr. 10 11 gr.
GASTRO-INTESTINAL IRR	ITAN TS.
Colchicum, radix 2-8 gr. "semen 1-5 gr. Creosotum ½-2 111. Digitalis ½-3 gr. Salicylic acid 5-60 gr. Scilla 1-5 gr. Strophanthus, tr. 2-10 11.	Sometimes in medicinal doses.
Alum	Useful as emetics.
Acidum carbolicum	

Guaiaci lignum	strongly.
Oleum tiglii	··· 100 10 gr.
Volatile oils. 4 Alcohol. Arnica, flores 10-20 gr. " radix 5-20 gr. Arsenic \$\frac{1}{10}\$ \ \ \frac{1}{10}\$ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	In large doses.

ALKALOIDS: THEIR SOURCES AND SALTS.

ACONITUM (monkshood):
 Aconitina
AMANITA MUSCARIA (fly fungus):
1 Muscarina 1-2 gr.
ASAGREA OFFICINALIS:
Veratrina (a mixture of alkaloids) 10 10 gr.
BELLADONNA (deadly nightshade):
1 Atropina 1 1 0 60 gr. Atropinæ sulphas 1 1 0 60 gr. Derived from atropina:
 IIomatropinæ hydrobromas 110 10 gr. Belladonina, probably identical with hyoscyamina.

CINCHONA:

I Quinina	I-20 gr.
Quininæ bisulphas	I-20 gr.
" hydrobromas	I-20 gr.
" valerianas	I-30 gr.
" sulphas	I-20 gr.
, "hydrochloras	I-20 gr.
2 Quinidina.	
Quinidinæ sulphas	I-20 gr.
3 Cinchonina	_
Cinchoninæ sulphas	I-30 gr.
4 Cinchonidina.	
Cinchonidinæ sulphas	1-30 gr.
COCA:	
Cocaina.	
Cocainæ hydrochloras	l -2 gr.
COLCHICUM:	
I Colchicina	l 1 gr.
Changed by acids into	
2 Colchiceina	0 64 gr.
DUDOICIA MYODODOIDEC.	

DUBOISIA MYOPOROIDES:

I Duboisina, probably identical with hyoscina. Duboisinæ sulphas, in discs containing tolog gr., to dilate pupil.

ERGOTA (secale cornutum, secale cereale):

- I Ergotina.
- 2 Ecbolina.
- 3 Ergotinina.
- 4 Cornutina.
- 5 Sphacelic acid (believed to contract the bloodvessels).

GELSEMIUM (yellow jasmine):

GRANATUM (pomegranate):

I Pelletierina.

Pelletierinæ tannas............ 8-24 gr. (in water 1 oz.: brisk cathartic.)

2 Three allied alkaloids.

HYDRASTIS (golden seal):

1 Hydrastina.

Derived from hydrastina:

Hydrastinina.

Hydrastininæ hydrochloras 1 gr.

```
ALKALOIDS: THEIR SOURCES AND SALTS.
                                                   140
    2 Berberina (also found in many plants),
                  dose { stomachic and tonic, \( \frac{1}{2} - \text{I} \) gr. antiperiodic...... 8-15 gr.
HYOSCYAMUS (henbane):
    1 Hyoscyamina.
         Hyoscyaminæ sulphas..... 1 50 80 gr.
                       hydrobromas.... \frac{1}{60} gr.
    2 Hyoscina (probably identical with duboisina).
         Hyoscinæ hydrobromas..... 180-180 gr.
    3 Scopolamina.
                             120
IPECACUANHA:
    Emetina (Merck's, dose 

emetic... 1 8 gr. expectorant, 1 8 gr.)
LOBELIA:
    Lobelina.
         Lobelinæ sulphas (Merck's, dose I gr. per day,
           gradually increasing to 3-8 gr. per day).
NUX VOMICA (poison nut, dog button):
    I Strychnina..... for no gr.
        Strychninæ sulphas..... \frac{1}{60-18} gr.
    2 Brucina.
    Strychnina is contained in other plants of the
      natural order Loganiacea.
```

150	ALKALOIDS: THEIR SOURCES AND SALTS.
OPIU .	M: •
1	Morphina.
	" acetas } 1 2 gr.
	Morphinæ sulphas
Der	ived from Morphina:
	Apomorphinæ hydrochloras,
	$dose \begin{cases} hypodermically, \frac{1}{10} & gr. \\ by mouth & \frac{1}{10} & 10 gr. \end{cases}$
2	Codeina ½-2 gr.
	Codeinæ phosphas 2 gr.
3	Thebaina.
4	Narceina ½ gr.
PHYS	SOSTIGMA (Calabar bean):
	hysostigmina = Eserine.
	Physostigminæ salicylas 100 80 gr.
	sulphas 100 80 gr.
PILO	CARPUS (jaborandi):
Pi	locarpina.
	Pilocarpinæ hydrochloras 1 gr.
SCOP	ARIUS (broom corn):
	parteina.
•	Sparteinæ sulphas

STRAMONIUM (thorn apple, Jamestown weed):

- I Daturina, a mixture of hyoscyamina and atropina.
- 2 Scopolamina.

-TABACUTI:

Nicotina.

VERATRUM VIRIDE (American hellebore):

- I Jervina.
- 2 Pseudo-jervina.
- 3 Veratroidina.

Pilocarpine Nicotine Comine

ACTIVE PRINCIPLES OF DRUGS OTHER THAN ALKALOIDS.

ADONI	S VERNALIS:
Ado	onidin, glucos de
ALOES	5 1 -10 gr.
Aloi	in, neutral principle $\frac{1}{2}$ gr.
ANAM	IRTA PANICULATA
Picz	IRTA PANICULATA. otoxin, neutral principle from seed, 110 6 gr
	IUM (male fern, filix-mas) \frac{1}{2}-1\frac{1}{2} dr. cic Acid, said to be active principle (Merck's, \frac{1}{2}-15 gr.)
BENZO	in:
Acid	lum benzoicum 5-15 gr.
CANTH	IARIS (Spanish fly):
	tharidin

ACTIVE PRINCIPLES OTHER THAN ALKALOIDS 15
CASSIA FISTULA (purging cassia) 1-8 dr
Cathartic Acid, glucoside, purgative principle
Also contained in senna,
Allo Contained in Delinia
CAFFEA ARABICA (coffee). 1
THEA SINENSIS (tea).
Caffeina, feebly basic, proximate principle, 1-5 gr
" citrata 2-8 gr
" effervescens 1-4 dr
COLOCVNTHIS (I'M - I - I - I - I
COLOCYNTHIS (bitter calumba) 2-8 gr Colocynthin, glucoside (Merck's, \(\frac{1}{6} - \frac{2}{8} \) gr.)
Colocynthin, glucoside (Merck s, 4-# gr.)
CONVALLARIA (lily of the valley):
Convallamarin, glucoside (Merck's, #-1 gr.)
CURARE (woorara, urari) gr
Curarina (Merck's: injection, 10-18 gr.)
CUSSO (brayera, kousso):
Kousin or kosin, neutral principle 20-40 gr
DIGITALIS (foxglove) 1-3 gr
The glucosides:
Digitoxin, the most active, poisonous,
cumulative (Merck's, 100 125 gr.)
Digitalin (Merck's; "German"; 1 1 gr)

Digitaleinsubcutaneously 100 gr.
Digitonin, probably identical with sa-
ponin of senega, cardiac depres-
sant 300 100 gr.
200 100 8
ELATERIUM (squirting cucumber):
Elaterin, neutral principle $\frac{1}{20-12}$ gr.
EUCALYPTUS ½-2 dr.
Eucalyptol, neutral body from oleum
eucalypti 5-30 ¶.
GALLA (nutgall):
Acidum tannicum
Acidum gallicum 5-20 gr.
GLYCYRRHIZA (licorice root):
Glycyrrhizin, glucoside 5-15 gr.
Glycyrrhizinum ammoniatum (official) 5-15 gr.
GOA POWDER (Araroba powder):
(Deposited in wood of andira araroba.)
Chrysarobinum (Rhein, chrysophan, chrysophanic
acid). Also from rheum.
GUARANA (Brazilian cocoa) 1-1 dr.
Guararina, identical with caffeina.

OTHER	THAN	ALKALOIDS.

HUMULUS (hops):
Lupulin, a glandular powder 5-30 gr.
Ext. fl 5-30 M.
Oleoresina 1-5 gr.
IRIS VERSICOLOR (blue flag).
Iridin, resin
JALAPA 5-20 gr.
Jalapurgin (convolvulin), glucoside. Most active ingredient.
Jalapin, resinous substance, probably identical with scammonin.
MENTHA PIPERITA (peppermint):
Menthol, a stearopten ½-2 gr.
PIPER (black pepper) 5-20 gr.
Piperinum, neutral principle 1-10 gr.
Also from others of order Piperacea.
PODOPHYLLUM (May apple, mandrake), 5-20 gr.
Podophyllin, official resin 1 gr.
Podophyllotoxin, resinous principle (Merck's,
1 - k gr.)
Saponin. See DIGITALIS and SENEGA.
QUASSIA:
Quassin, bitter principle 1 gr.

156 ACTIVE PRINC	CIPLES OTHER THAN ALKA	LOIDS.
Santonin, neut	tral principledose { child adult	15-60 gr. 1-1 gr. 1-5 gr.
SCAMMONIUM.		5-15 gr.
Scammonin, r	resinous substance, probabl	y identi-
Scillitoxin (scill tive principle Scillipierin, b neously, 1-1	9.1	
	onin, glucoside, acts like d	
	, glucoside, purgative princ	
STROPHANTHU		
	glucoside (Merck's, store 1 to 1 t	
OLEUM THYMI	Dan - In the last	
	enol (a stearopten in B. P.)	

Thymol, a phenol (a stearopten in B. P.) 1-15 gr. Also from volatile oils of monarda punctata and carum ajowan.

DRUGS WITH MAXIMUM DOSE ONE-TENTH GRAIN OR LESS.

I	Acidum arsenosum	1 1 1	gr.
2	Aconitina	200	gr.
-	(with caution)		
3	Apomorphinæ hydrochloras,		
	dose { by mouth injection	20 10	gr.
	dose (injection	25 6	gr.
4	Atropina	120 60	gr.
5	Atropinæ sulphas	120 20	gr.
6	Auri et sodii chloridum	10 10	gr.
7	Auri et sodii chloridum	100	gr.
	(injection.)	***	
8	Digitalin (Merck's "German")	64 33	gr.
9.	Digitopin	300 100	gr.
10	Digitonin Elaterinum Hydrargyri chloridum corrosivum	30 13	gr.
II	Hydrargyri chloridum corrosivum	1 1	gr.
12	" iodidum rubrum	1 1	
13	Hyoscinæ hydrobromas	40 40	-
-	Hyoscyaminæ "		
15	" sulphas	40 40	-
+3	sarbagair :	00 80	8.

16 Nitroglycer (containe	inum d in spr. glon	oini.)	100 8
17 Phosphorus	(in pill)		$\frac{1}{100} - \frac{1}{2}$
18 Physostigm	inæ salicylas.		100 8
19 "			
20 Sodii arsena	s		80 1
21 Strychnina			60 2
22 Strychninæ	sulphas		60 1
Sines	amine		120

DRUGS KNOWN BY COMMON AND OTHER NAMES.

Adhesive plaster—Emplastrum resine. Ammonia, baker's—Ammonii carbonas, 2-15 gr.

Basham's mixture—Liquor ferri et ammonii acetatis,
 1-4 dr.

Basilicon ointment-Ceratum resinæ,

Battley's solution—Liquor opii sedativus (not official), containing 50%+ of tr. opii.

Black draught-Infusum sennæ compositum, 1-2 oz.

-Black drop-Acetum opii (10%), 3-20 ₪.

Also applied to Acetum opii crocatum (40%).

-Blaud's pill-Pilulæ ferri carbonatis, 1-2 pills.

-Bleaching powder-Calx chlorata.

←Blue mass—Massa hydrargyri (33%).

ointment—Unguentum hydrargyri

" pill—Massa hydrargyri.

vitriol Cupri sulphas, astringent 1 gr. cmetic... 2-20 gr.

- Brandy-Spiritus vini gallici (39-47%).

Broom corn-Scoparius, 1-1 dr.

Brown mixture—Mistura glycyrrhizæ composita, 1-4 dr.

Burnett's fluid-Solution of zinc chloride, sp. gr. 2.0.

Cacao butter-Oleum theobromatis.

Cade, oil of—Oleum cadini (Oleum juniperi empyreumaticum).

Calabar bean—Physostigma, 1-2 gr. of powdered beans.

Calomel-Hydrargyri chloridum mite, 10-10 gr

Carron oil-Linimentum calcis.

Castile soap (hard soap)-Sapo.

Castor oil-Oleum ricini, 1-2 oz.

Cayenne pepper-Capsicum, 1-8 gr.

Chalybeate pill—Pilulæ ferri carbonatis, 1-2 pills.
Cherry, wild—Prunus Virginiana, 1-1 dr.

- Chloric ether—Spiritus chloroformi, 1-1 dr.

Cod-liver oil—Oleum morrhuæ (Oleum jecoris aselli),

Cold cream-Unguentum aquæ rosæ.

Common salt-Sodii chloridum, 5-60 gr.

Condy's fluid-Potassii permanganas, 8 gr.; water, 1

Copperas (green vitriol)—Ferri sulphas (impure).

Corrosive sublimate—Hydrargyri chloridum corrosivum, 20 70 gr.

Coster's paste—Iodine, 2 dr.; oleum picis liquidæ,

Cotton, absorbent—Gossypium purificatum.
Couch grass—Triticum, 1-8 dr.
Court plaster—Emplastrum ichthyocollæ.
Croton oil—Oleum tiglii, ½-2 M.
Culver's root—Leptandra, ¼-1 dr.

Dandelion-Taraxacum, 1-3 dr.

- Deadly nightshade—Belladonna (folia), 1-5 gr.
- Diachylon plaster—Emplastrum plumbi.

Dinner pill—Pilulæ aloes et mastiches, 1-3 pills. Dog button (poison nut)—Nux vomica, 1-4 gr.

- ←Donovan's solution—Liquor arseni et hydrargyri iodidi (1≴ of each), 1-10 Щ.
- ∠ Dover's powder—Pulvis opii et ipecacuanhæ (10% of each), 3-15 gr.

Dutch liquid-Æthylenum bichloridum.

Epsom's salt—Magnesii sulphas, 1-1 oz. Eserine—Physostigmina.

Fern, male (filix-mas)—Aspidium, 1-11 dr.

Fleming's tincture—A tincture of aconite twice as strong as official tincture, ½-2 M.

Foxglove-Digitalis, 1-3 gr.

→Fowler's solution—Liquor potassii arsenitis (τ\$), 2-

Friar's balsam—Tinctura benzoini composita, 1-1 dr. Fusel oil—Alcohol amylicum.

- Glauber's salt-Sodii sulphas, 1-8 dr.
- Goulard's cerate-Ceratum plumbi subacetatis.
- extract—Liquor plumbi subacetatis.
- ← Gray powder-Hydrargyrum cum creta (38%), 1-10 gr.
- ∠ Green vitriol—Ferri sulphas (impure).
- Gregory's powder—Pulvis rhei compositus, 1-1 dr.
 Griffith's mixture—Mistura ferri composita, 2-4 dr.
- Gypsum (plaster of Paris)—Calcii sulphas exsiccatus.
- Hartshorn-Ammonii carbonas, 2-15 gr.
- ► Haschisch-Cannabis Indica, 2-5 gr.

Hellebore, American-Veratrum viride, 1-5 gr.

- Hemlock-Conium, 2-5 gr.
- ✓ Hemp, Indian—Cannabis Indica, 2-5 gr.
 - ' Canadian-Apocynum,

dose { emetic ... 10-20 gr. alterative, 2-5 gr.

- Henbane-Hyoscyamus, 5-15 gr.
- ← Hive syrup—Syrupus scillæ compositus, 5-30 M.
- ← Hoffman's anodyne—Spiritus ætheris compositus, 5-60 ¶.

Jaborandi—Pilocarpus, 5-60 gr.

James' powder—Pulvis antimonialis, 3-15 gr.

- ← Jamestown weed ("Jimson" weed)—Stramonium, (seed) 1-3 gr., (leaves) 1-5 gr. Jasmine, yellow—Gelsemium, 5-10 gr.
- Labarraque's solution—Liquor sodæ chloratæ, ½—1 dr.
 "Lady Webster" pill—see Dinner pill.
- Laudanum—Tinctura opii (10%), 3-20 ℍ. Licorice, compound powder—Pulvis glycyrrhizæ compositus, ½-2 dr.
- Lugol's solution—Liquor iodi compositus (5%), 1-10
- Lunar caustic-Argenti nitras, 1-1 gr. in pill.

Mandrake Podophyllum, 5-20 gr.

- Mercurial ointment-Unguentum hydrargyri (50%).
- Mercury biniodide—Hydrargyri iodidum rubrum, 10-10 gr.
- Mercury protoiodide—Hydrargyri iodidum flavum, a-1 gr.
- Mindererus, spirit of—Liquor ammonii acetatis, 2-8 dr.
- ∠ Monsel's solution—Liquor ferri subsulphatis, 2-10 M.
- Morphia, U. S. P. solution of—Morphine sulphas, 1 gr.; water, 1 oz.

- Nitre (saltpetre)-Potassii nitras, 5-20 gr.
- " sweet spirit of—Spiritus ætheris nitrosi, ½-2 dr. Norwood's tincture—Tinctura veratri viridis, 2-10 π.
- Oil of vitriol—Acidum sulphuricum.
 Opodeldock—Linimentum saponis.
- Paregoric—Tinctura opii camphorata (2 gr. to 1 oz.),
 1-1 dr.
- ► Pearson's solution—Liquor sodii arsenatis (♣), 1-10 M.

 Pearson's solution is really one-tenth as strong as
 the official liquor sodii arsenatis. 2 9
- Plaster of Paris-Calcii sulphas exsiccatus.
- ← Plummer's pills Pilulæ antimonii compositæ, 1-3 pills.
 Poison nut—Nux vomica, 1-4 gr.
 Pomegranate—Granatum, ½-1½ dr.
- Poppy-Opium, 1-2 gr.
- Port wine-Vinum portense.
- ► Precipitate, red—Hydrargyri oxidum rubrum, 1-1 gr.
- white-Hydrargyrum ammoniatum.
- Prussic acid—Acidum hydrocyanicum,
 Pulvis purgans—Pulvis jalapæ compositus, 1-1 dr.
 Pumpkin seed—Pepo, 1-3 oz.
- Quevenne's iron-Ferrum reductum, 1-5 gr.
- 4 Quicksilver-Hydrargyrum.

4 Rochelle salt-Potassii et sodii tartras,

Sacred bark—Rhamnus purshiana (cascara sagrada).

- Sal ammoniac—Ammonii chloridum, 1-30 gr.
- " soda Sodii earbonas, 5-15 gr.
- " volatile—Spiritus ammoniæ aromaticus, 1-2 dr.
- Salts of tartar-Potassii carbonas, 5-30 gr.
- Seidlitz powder—Pulvis effervescens compositus, I powder.
- ← Sherry—Vinum xericum.
- Soda, baking—Sodii bicarbonas, 5-60 gr.
 Spanish fly—Cantharis.
- Spirit of nitroglycerine—Spiritus glonoini (1%), 1-3 π.
- ← Sugar of lead—Plumbi acetas, ½-5 gr.
- Sydenham's laudanum-Vinum opii (10%), 3-20 Π.
- Tartar emetic—Antimonii et potassii tartras,
- Tartar, cream of-Potassii bitartras,

dose { diuretic 20-60 gr. purgative
$$\frac{1}{2}$$
-4 dr.

- ← Thornapple seed—Stramonii semen, 1-3 gr.
- Tully's powder—Pulvis morphimæ compositus (1 gr. to 1 dr.), 5-15 gr.
- Turpeth mineral—Hydrargyri subsulphas flavus,
 2-4 gr.

166 DRUGS KNOWN BY COMMON AND OTHER NAMES.

- ← Vallet's mass—Massa ferri carbonatus, 1-5 gr.
- Vienna paste Potassa cum calce.

Warburg's tincture—Tinctura antiperiodica (2% quinine), 1-4 dr.

Warming plaster-Emplastrum picis cantharidatum.

Washing soda—Sodii carbonas, 5-15 gr.

Wild cherry-Prunus Virginiana, 1-1 dr.

Witchhazel-Hamamelis.

Wintergreen, oil of-Oleum gaultheriæ, 1-5 M.

Catanhal Janudice (Loomis

High rectal imig. 65° 7

Loca a day. 3. 4 gt.

Na. HPOY 12 & before health

Retinated will diet.

For acute Bronchitis

Redeinae gr. 7111

Acidi Hydrocyanii Kiluti M MXII

Annonii Chloridi Z 17%

Syrupi pruni Virginianae gs. ad ZI

DOSAGE IN CHILDREN.

Young's rule:

$$\frac{age}{age + 12} = x = fraction of adult dose.$$

Example:

$$\frac{2}{2+12}=\frac{1}{7}.$$

Cowling's:

$$\frac{\text{age} + 1}{24} = x.$$

Example:

$$\frac{2+1}{24}=\frac{1}{8}.$$

RULES FOR INCOMPATIBILITY OF DRUGS.

- Mineral acids are incompatible with the alkalies, their carbonates, and lead.
- Metallic salts are precipitated by alkalies and tannic acid.
- III. The alkaloids are incompatible with tannic acid and caustic alkalies.
- IV. All drugs are incompatible with their antidotes and chemical tests.

Chologogue

Noi Sal : gr. 11]

Ext. coloh. noot : 12-1

Tellio Borio : 2-3

Res. Podophighi : 12-4

J. i. d.

AN EASY METHOD OF WRITING PRESCRIPTIONS.

It may be assumed, for the purpose of writing prescriptions, that there are 15 doses of a teaspoonful each in a 2-ounce mixture, 30 in a 4-ounce mixture, and 60 in an 8-ounce mixture. Only in the case of dangerous drugs is a more accurate estimation necessary.

In a 4-ounce mixture, then, with a teaspoonful dose, each dose will contain $\frac{1}{30}$ of the total amount of any drug which may be in solution or uniform suspension.

In the case of drugs with a usual dose of about 5 gr. or M, I drachm may be taken as the basis of calculation.

If I drachm of a drug be added to a 4-ounce mixture, each teaspoonful will contain $\frac{1}{10}$ of a drachm, or 2 gr. or η .

Taking 2, then, as a unit, it is only necessary to find the multiple of 2 which will give the desired dose, and this will represent the number of drachms to be put into the prescription.

To take an example:

Encalyptol m

AN EASY METHOD OF WRITING PRESCRIPTIONS, 160

In a 2-ounce mixture, each teaspoonful will contain $\frac{1}{15}$ of a drachm, or 4 gr. or \mathbb{N} .

In an 8-ounce mixture, each teaspoonful will contain $\frac{1}{6}$ of a drachm, or I gr. or M.

From the above statements the following rule may be formulated:

Divide 60 (one drachm) by the number of doses in the prescription, and multiply the result by the numeral necessary to give the desired dose. This numeral will represent the number of drachms to be used.

In the case of drugs with a maximum dose of less than a grain, I grain instead of I drachm may be taken as the basis of calculation. Thus, if I grain be added to a 4-ounce mixture with a teaspoonful dose, each dose will contain $\frac{1}{100}$ of a grain.

1) Marphine - 4 gr. by hypo.

2) Wine of Colchieum Noot - 20 gth (10m) in

Repeat in 4 & lathird deserming not be my

3) Baline - large dose - in the manual

Discontinue use of colchieum when col

pairs begins foot in cotton for

y) What mp foot in cotton for


protection a mint lust.

Hairo fift also.

Pel of Pul iperat a 12 gr. Jan.

a. & C. Mulime : 4 mars Heite 219 Va. B. C. Summer Falladoura comphor as Guy riples Sqlill
Bigitales
calonel an 19r
6x agosegamo 12. Buladone Places would chalera mixture In , april " enpaire"
" plain
" exmellerier
" " menter pipuitar aa a do me has the D Los Jake marins and big of annual and to E. Eng. punder the righ ratefast except a cut of est

3 maturent of Syphilis (Hare) Protionide of Hg in 4 gr. tablet tri Begin with '3 tablets a day and & (Hysolicylate 1/2 gr amanink For salivation In bellasonne 10 m enny 3 h. they muth aufter + alkeline (Somia) month wash KCLO: 1 track, to the glass In. murch For chanic broughitis (Have) Paregnic Squill For malingeners Jr. green sout .. Valarian .. asafetida Give me on two track, in Viely. For tapensony (Dr. Coleman) 9.2. of the alkaloids of grandim. missing to 12/2 for 13 bottle. Hore 13 for which



U122 Coleman, W. 108659 C69 A syllabus of materia 1899 medica. DATE DUE NAME

